

Annual Data Summary
HAWAII VOLCANOES NATIONAL PARK
1998
National Park Service
Gaseous Air Pollutant Monitoring Network



AIR RESOURCES DIVISION
RESEARCH AND MONITORING BRANCH
12795 West Alameda Parkway
P.O. Box 25287
Lakewood, Colorado 80225
Telephone: (303) 969-2820
Fax: (303) 969-2822

This Annual Data Summary was prepared under NPS Contract CX-1270-96-007 by:

Air Resource Specialists, Inc.
1901 Sharp Point Drive, Suite E
Fort Collins, Colorado 80525
Telephone: (970) 484-7941
Fax: (970) 484-3423

For additional copies of this report or reports for other NPS units, contact:

National Park Service Air Resources Division
Information Management Center
c/o Air Resource Specialists, Inc.
1909 Sharp Point Drive, Suite E
Fort Collins, Colorado 80525
Telephone: (970) 484-7941
Fax: (970) 484-3423
E-Mail: AIR-IMC@AIR-RESOURCE.COM

or

National Park Service
Air Resources Division
P.O. Box 25287
Lakewood, Colorado 80225-02587
Telephone: (303) 969-2130
E-Mail: AQ_INFO@AQD.NPS.GOV

ACKNOWLEDGEMENTS

The National Park Service Air Resources Division (ARD) recognizes the level of effort required by individual park units, site operators, auditors, cooperating state and local agencies, and ARD contractors. ARD sincerely appreciates the contributions of all participants in assisting with the collection, validation, and reporting of these air quality and meteorological data.

At Hawaii Volcanoes National Park, the ARD specifically recognizes Tamar Elias for performing the technical and administrative skills required to help produce the data presented within this report.

TABLE OF CONTENTS

<u>Section</u>		<u>Page</u>
1.0	INTRODUCTION	1-1
1.1	The National Park Service Gaseous Pollutant Monitoring Network	1-1
1.2	Hawaii Volcanoes National Park	1-3
2.0	DATA SUMMARY	2-1
2.1	Overview	2-1
2.2	Sulfur Dioxide Data Summary	2-7
2.3	Supplemental Sulfur Dioxide Data Summary	2-19
2.4	Meteorological Data Summary	2-27
3.0	NATIONAL PARK SERVICE AIR RESOURCES DIVISION DATA SOURCES	3-1
3.1	Guide to Attached Data Disks	3-1
3.2	Other Sources for Retrieving National Park Service Gaseous Pollutant Data	3-1
4.0	GLOSSARY	4-1
4.1	Definitions and Computational Procedures for National Park Service Quick Look Annual Summary Statistics Table	4-1
4.2	Air Quality Glossary	4-4

LIST OF FIGURES AND TABLES

	<u>Page</u>
Map of National Park Service Gaseous Pollutant Monitoring Network Ozone and Sulfur Dioxide Monitoring Sites	1-2
Site Specifications	1-4
Data Collection Statistics	2-2
First Quarter Data Stackplot	2-3
Second Quarter Data Stackplot	2-4
Third Quarter Data Stackplot	2-5
Fourth Quarter Data Stackplot	2-6

LIST OF FIGURES AND TABLES (Continued)

	<u>Page</u>
Sulfur Dioxide Daily 1-Hour Maximum Concentrations and National Ambient Air Quality Standards Comparison	2-8
Sulfur Dioxide Standards, 1-Hour Maximum	2-9
Sulfur Dioxide Standards, Maximum of 3-Hour Block Average	2-10
Sulfur Dioxide Standards, 24-Hour Block Average	2-11
Sulfur Dioxide Yearly Frequency Distribution	2-12
Maximum Sulfur Dioxide Concentration Comparisons for Three Years	2-13
Quarterly Diurnal Sulfur Dioxide Plots	2-14
Annual Diurnal Sulfur Dioxide Plots	2-15
Quarterly Sulfur Dioxide Pollutant Roses	2-16
Annual Sulfur Dioxide Pollutant Roses	2-17
Sulfur Dioxide Precision Check Data Summary	2-18
Supplemental Sulfur Dioxide Daily 1-Hour Maximum Concentrations and National Ambient Air Quality Standards Comparison	2-20
Supplemental Sulfur Dioxide Three Year Comparison	2-21
Quarterly Diurnal Supplemental Sulfur Dioxide Plots	2-22
Annual Diurnal Supplemental Sulfur Dioxide Plots	2-23
Quarterly Supplemental Sulfur Dioxide Pollutant Roses	2-24
Annual Supplemental Sulfur Dioxide Pollutant Roses	2-25
Supplemental Sulfur Dioxide Precision Check Data Summary	2-26
Summary of Selected Meteorological Data	2-28
Quarterly Wind Roses	2-29
Annual Wind Rose	2-30
Data Disk Contents	3-2
NPS IMC and AIRS Invalid Data Codes	3-3

1.0 INTRODUCTION

1.1 THE NATIONAL PARK SERVICE GASEOUS POLLUTANT MONITORING NETWORK

Gaseous air pollutants, including ozone and sulfur dioxide, are of concern to the National Park Service (NPS). Pollutants like these can affect park unit biological resources as well as the health of park unit residents and visitors. The NPS established a gaseous pollutant monitoring program for several pollutants linked to effects on NPS resources. This program was designed to meet certain resource management objectives.

The primary objective of this monitoring program is to establish the status and trends of park unit air quality conditions and to determine if a park unit is exceeding the National Ambient Air Quality Standards established by the U.S. Environmental Protection Agency (EPA) to protect public health and welfare. In addition, such monitoring is designed to detect changes or trends in pollution levels over time. A monitoring station may also be established if there is documented biological injury due to air pollution in a park unit. Information on ambient air pollution levels is an important part of research on effects of air pollutants on NPS resources, and can help confirm suspected causes of observed effects.

Other monitoring objectives call for the collection of data to support the National Park Service's required involvement in both the development of state air quality control plans, and the evaluation of permit applications for new or expanding air pollution sources wishing to locate near park units. The Clean Air Act gives federal land managers and superintendents an affirmative responsibility to protect air quality related values in Class I areas and to assess whether new sources will have an adverse impact on park unit resources and values. Information on air quality levels in NPS units can also be used to evaluate the performance of atmospheric models that simulate how pollutants are transported into park units and predict impacts on the park unit caused by air pollution sources.

The National Park Service Gaseous Pollutant Monitoring Network site locations and measured parameters collected in this reporting year are shown on the map on the following page. During this reporting period, 40 monitoring sites in 35 units of the National Park System had some combination of ozone, sulfur dioxide, meteorological, and CASTNet dry deposition monitoring. Monitoring methods and quality assurance procedures used in the national park network meet the applicable 40 CFR Part 58 EPA requirements. This allows for the direct comparison of NPS collected data with that collected by the EPA, and state and local air pollution control agencies. Data collected by this network are incorporated in the EPA Aerometric Information Retrieval System (AIRS) database which is a national database of all air quality data collected throughout the country. These data are also stored in the NPS Air Resources Division's Information Management Center (IMC) that allows for easy access and analysis of data.

This report includes a variety of data summaries for data collected at an individual monitoring site at a national park unit during this reporting period. These summaries highlight the average range and frequency of the data collected during the year. A PC-compatible diskette containing a digital copy of all data collected during the year and data summary products included in this report is available. Individual reports are generated for each site where monitoring was conducted in the national park network.

**NATIONAL PARK SERVICE
GASEOUS POLLUTANT MONITORING NETWORK**

1998 Ozone and Sulfur Dioxide Monitoring Sites



1.2 HAWAII VOLCANOES NATIONAL PARK

Hawaii Volcanoes National Park, a Class I area, is located in the southeastern portion of the island of Hawaii. The island is situated at the southeastern end of the Hawaiian chain and is the largest and most recently formed. Its location and site specifications are presented on the following page.

The park was established by Congress in 1916 to conserve the volcanic features, endemic Hawaiian ecosystems, Hawaiian cultural and archeological remains, and inherent scenic values for visitor enjoyment and appreciation and for their scientific and historic values. In 1987, Congress designated 123 thousand of the park's 229 thousand acres as a wilderness area. Hawaii Volcanoes National Park was designated a Biosphere Reserve in 1980, and a World Heritage Site in 1987.


Hawaii Volcanoes National Park contains within its boundaries two of the most active volcanoes in the world, Mauna Loa and Kilauea, along with numerous historical, archeological, and biological resources.

Kilauea and Mauna Loa volcanoes are the dominant features of the park. Surrounded by recent lava flow materials and unique endemic plant communities, these volcanoes are dynamic landforms where new lava flows can drastically change the landscape. Recent eruptions have covered many acres with lava up to 300 feet deep, created new land where flows enter the ocean, and built up a new mountain where none existed before. These two volcanoes are the most studied and best understood volcanoes in the world because of the favorable opportunities afforded for fundamental and detailed research which are not duplicated or even approached in any other part of the world.

Hawaii flora is remarkable for its extraordinary degree of endemism. Ninety-five percent of the native species are found only in these islands. Although park boundaries were drawn primarily to include recent lava flows and the summits of Kilauea and Mauna Loa volcanoes, there is within the park, albeit limited in extent, a rich diversity of species and vegetation types. Diverse natural vegetation zones include tropical rain forest, woodlands, shrublands, grasslands, and alpine tundra.

Birds are the most important aspect of the park's wildlife and include endemic species of honeycreeper, owl, thrush, hawk, and flycatcher. No fish are found within the park. The Hawaiian bat is the only native land mammal. However, feral animals, such as goats and pigs, have done great damage to the native plant and animal communities.

The park's cultural and historical resources include Hawaiian religious sites, prehistoric and historic trails and routes and relatively modern historic structures.



SITE IDENTIFICATION		MAP INFORMATION	
Site Abbreviation:	HAVO	Mean Elevation:	1215 m
AIRS ID NO.:	15-001-0005	Longitude:	155° 15' 40"W
INSTRUMENTATION		Latitude:	19° 26' 01"N
		UTM Zone:	05
SO ₂ Analyzer	Dew Point	Easting:	262577 m
Wind Speed	Temperature	Northing:	2150392 m
Wind Direction	Solar Radiation	Map Reference:	Hawaii
Relative Humidity	Precipitation		NE 5-1,5
			1975
			1:250,000

2.0 DATA SUMMARY

2.1 OVERVIEW

Based on the site specifications during this annual reporting period, data summaries and statistics are provided in this section.

Data Collection Statistics
Hawaii Volcanoes National Park

Final Data

01/01/98 - 12/31/98

Parameter	Par Code	Data Recovery			Valid Data	
		No. Possible	No. Collected	% Collected	No. Valid	% Valid
Sulfur Dioxide Analyzer	SO2	8760	8334	95.1	8129	92.8
SO2 Additional Analyzer	SO2ADD	8760	8399	95.9	6873	78.5
Scalar Wind Speed	SWS	8760	8739	99.8	7595	86.7
Vector Wind Speed	VWS	8760	8734	99.7	5919	67.6
Vector Wind Direction	VWD	8760	8734	99.7	7798	89.0
Ambient Temperature (aspirated)	TMP	8760	8719	99.5	8591	98.1
Relative Humidity	RH	6600	6599	100.0	4611	69.9
Dew Point	DPT	1967	1967	100.0		
Precipitation	RNF	8760	8730	99.7	8730	99.7
Solar Radiation	SOL	8760	8741	99.8	8741	99.8

Notes: All statistics are for hourly averages.

The number collected does not include normal maintenance or events beyond the control of the network.

The percent valid is calculated against the number possible.

Automatic zeros and spans are performed daily on most ambient gas analyzers, therefore, no ambient data can be collected during this time. As a result, the maximum percent valid for ambient gas data typically can not be greater than 95.8.

NPS Performance Goals:

Quarterly Criteria:


100% of sites, >= 85% valid data capture
 90% of sites, >= 90% valid data capture
 80% of sites, >= 95% valid data capture

Monthly Criteria:

100% of sites, >= 60% valid data capture
 90% of sites, >= 75% valid data capture
 80% of sites, >= 85% valid data capture

* The validity of dew point and relative humidity data is currently under investigation.

Hawaii Volcanoes National Park




Final Validation

First Quarter 1998

havo-vc.stk - havo98.dat 08-23-1999

Hawaii Volcanoes National Park




Final Validation

Second Quarter 1998

havo-vc.stk - havo98.dat 08-23-1999

Hawaii Volcanoes National Park




Final Validation

Third Quarter 1998

havo-vc.stk - havo98.dat 08-23-1999

Hawaii Volcanoes National Park



Final Validation

Fourth Quarter 1998

havo-vc.stk - havo98.dat 08-23-1999

2.2 SULFUR DIOXIDE DATA SUMMARY

Sulfur Dioxide
 Five Highest Daily 1-Hour Averages, 3-Hour Block Averages,
 and 24-Hour Block Averages
 Hawaii Volcanoes National Park

Final Data
 01/01/98 - 12/31/98

Rank	Date	Hour	Concentration (ppb)
Highest Daily 1-Hour Averages			
1	01/15/98	12	952
2	11/11/98	10	912
3	11/08/98	15	903
4	08/11/98	8	871
5	11/12/98	9	852
Highest 3-Hour Block Averages			
1	11/11/98	9-11	649
2	01/15/98	12-14	648
3	11/11/98	6-8	567
4	11/05/98	9-11	533
5	01/16/98	0-2	492
Highest 24-Hour Block Averages			
1	11/11/98	00-23	302
2	01/14/98	00-23	258
3	01/15/98	00-23	191
4	08/11/98	00-23	157
5	11/08/98	00-23	157

Note: The primary and secondary ambient air standards for SO₂ are:
 Annual Primary Standard - 0.03 ppm; annual arithmetic mean not to be exceeded.
 (A value greater than .03 ppm, 34 ppb, or 80 ug/m³ exceeds the standard.)
 24-Hour Primary Standard - 0.14 ppm; not to be exceeded more than once per year.
 (A value greater than .14 ppm, 144 ppb, or 365 ug/m³ exceeds the standard.)
 3-Hour Secondary Standard - 0.5 ppm; not to be exceeded more than once per year.
 (A value greater than .5 ppm, 549 ppb, or 1300 ug/m³ exceeds the standard.)
 (40 CFR 50.4 and 50.5 with reference to Appendix A.)

Sulfur Dioxide Standards Report and
Daily Maximum 1-Hour Concentrations (ppm)
Hawaii Volcanoes National Park
01/01/98 - 12/31/98

Day	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec												
1	T	.658	S	.012	S	.003	W	.000	F	.000	M	.000	W	.399	S	.003	T	.002	T	.002	S	.001	T	
2	F	.001	M	.025	M	.000	T	.000	S	.000	T	.000	T	.002	S	.002	W	.002	F	.002	M	.001	W	
3	S	.224	T	.000	T	.000	F	.000	S	.000	W	.000	F	.002	M	.002	T	.002	S	.002	T	.001	T	
4	S	.168	W	.226	W	.000	S	.000	M	.000	T	.000	S	.002	T	.002	F	.002	S	.002	W	.002	F	
5	M	.132	T	.273	T	.000	S	.000	T	.000	F	.000	S	.002	W	.002	S	.002	M	.619	T	.002	S	
6	T	.013	F	.144	F	.000	M	.000	W	.000	S	.000	M	.002	T	.002	S	.002	T	.053	F	.001	S	
7	W	.003	S	.187	S	.000	T	.000	T	.000	S	.000	T	.002	F	.002	M	.002	W	.685	S	.001	M	
8	T	.201	S	.087	S	.000	W	.000	F	.000	M	.000	W	.002	S	.002	T	.002	T	.903	S	.657	T	
9	.223	F	.043	M	.034	M	.000	T	.000	S	.000	T	.000	T	.002	S	.002	W	.002	F	.003	M	.002	W
10	.578	S	.122	T	.000	T	.000	F	.000	S	.000	W	.000	F	.006	M	.002	T	.002	S	.586	T	.001	T
11	.334	S	.219	W	.000	W	.000	S	.000	M	.000	T	.000	S	.871	T	.002	F	.002	S	.912	W	.001	F
12	.009	M	.074	T	.005	T	.000	S	.000	T	.000	F	.000	S	.003	W	.002	S	.062	M	.852	T	.001	S
13	.629	T	.002	F	.462	F	.000	M	.000	W	.000	S	.000	M	.002	T	.002	S	.002	T	.001	F	.001	S
14	.844	W	.000	S	.446	S	.000	T	.000	T	.000	S	.000	T	.002	F	.002	M	.002	W	.002	S	.384	M
15	.952	T	.001	S	.055	S	.000	W	.000	F	.000	M	.000	W	.002	S	.002	T	.002	T	.002	S	.118	T
16	.642	F	.000	M	.009	M	.000	T	.000	S	.000	T	.000	T	.017	S	.002	W	.002	F	.002	M	.189	W
17	.063	S	.004	T	.000	T	.000	F	.000	S	.000	W	.000	F	.803	M	.002	T	.002	S	.001	T	.086	T
18	.002	S	.000	W	.000	W	.000	S	.000	M	.000	T	.000	S	.004	T	.002	F	.002	S	.001	W	.058	F
19	.000	M	.000	T	.023	T	.000	S	.000	T	.000	F	.000	S	.002	W	.002	S	.003	M	.001	T	.243	S
20	.102	T	.000	F	.000	F	.001	M	.000	W	.000	S	.000	M	.002	T	.002	S	.002	T	.002	F	.002	S
21	.204	W	.000	S	.088	S	.000	T	.000	T	.000	S	.002	T	.002	F	.002	M	.002	W	.002	S	.001	M
22	.070	T	.001	S	.000	S	.000	W	.000	F	.000	M	.002	W	.002	S	.002	T	.002	T	.002	S	.001	T
23	.238	F	.000	M	.000	M	.000	T	.000	S	.000	T	.002	T	.002	S	.002	W	.010	F	.002	M	.001	W
24	.146	S	.000	T	.005	T	.000	F	.000	S	.000	W	.002	F	.002	M	.002	T	.002	S	.001	T	.002	T
25	.026	S	.000	W	.008	W	.000	S	.000	M	.000	T	.005	S	.002	T	.002	F	.002	S	.001	W	.001	F
26	.596	M	.005	T	.003	T	.085	S	.000	T	.000	F	.002	S	.002	W	.002	S	.002	M	.001	T	.001	S
27	.388	T	.005	F	.003	F	.323	M	.000	W	.000	S	.002	M	.002	T	.002	S	.002	T	.002	F	.473	S
28	.001	W	.184	S	.003	S	.002	T	.000	T	.000	S	.002	T	.002	F	.002	M	.003	W	.002	S	.203	M
29	.141	T			.003	S	.000	W	.000	F	.000	M	.002	W	.002	S	.002	T	.002	T	.001	S	.281	T
30	.021	F			.003	M	.000	T	.000	S	.000	T	.002	T	.002	S	.002	W	.002	F	.001	M	.228	W
31	.001	S			.003	T			.000	S			.002	F	.002	M			.002	S			.152	T
Number	515	638	695	684	708	685	708	709	685	709	685	708												
Maximum	.952	.658	.462	.323	.000	.000	.005	.871	.003	.062	.912	.657												
Arith-Mean	.048	.012	.014	.004	.001	.001	.001	.012	.001	.002	.029	.019												
8129 Total Samples			.011 Arith-Mean					Statistics meet summary criteria																
92.8 % Possible			.059 Arith-StdDev					Primary standard of .030 ppm was met				Concentrations in parts per million (ppm)												

Sulfur Dioxide Standards Report and
Daily Maximum 24-Hour Concentrations (ppm)
Hawaii Volcanoes National Park
01/01/98 - 12/31/98

Day	Jan-94	Feb-94	Mar-94	Apr-94	May-94	Jun-94	Jul-94	Aug-94	Sep-94	Oct-94	Nov-94	Dec-94
1	T	.060 S	.002 S	.002 W	.001 F	.001 M	.001 W	.039 S	.001 T	.001 T	.001 S	.001 T
2	F	.001 M	.003 M	.001 T	.001 S	.001 T	.001 T	.002 S	.001 W	.002 F	.001 M	.001 W
3	S	.057 T	.001 T	.001 F	.001 S	.001 W	.001 F	.001 M	.001 T	.002 S	.001 T	.001 T
4	S	.032 W	.058 W	.001 S	.001 M	.001 T	.001 S	.002 T	.001 F	.002 S	.001 W	.001 F
5	M	.021 T	.057 T	.001 S	.001 T	.001 F	.001 S	.002 W	.001 S	.001 M	.101 T	.001 S
6	T	.002 F	.031 F	.001 M	.001 W	.001 S	.001 M	.002 T	.001 S	.001 T	.005 F	.001 S
7	W	.001 S	.022 S	.001 T	.001 T	.001 S	.001 T	.002 F	.001 M	.001 W	.049 S	.001 M
8	T	.043 S	.011 S	.001 W	.001 F	.001 M	.001 W	.002 S	.001 T	.001 T	.157 * S	.153 * T
9	F	.008 M	.003 M	.001 T	.001 S	.001 T	.001 T	.002 S	.002 W	.001 F	.002 M	.001 W
10	.121 S	.024 T	.001 T	.001 F	.001 S	.001 W	.001 F	.002 M	.002 T	.001 S	.085 T	.001 T
11	.059 S	.032 W	.001 W	.001 S	.001 M	.001 T	.001 S	.157 * T	.001 F	.001 S	.302 * W	.001 F
12	.002 M	.023 T	.001 T	.001 S	.001 T	.001 F	.001 S	.002 W	.001 S	.005 M	.137 T	.001 S
13	.062 T	.001 F	.094 F	.001 M	.001 W	.001 S	.001 M	.002 T	.002 S	.001 T	.001 F	.001 S
14	.258 * W	.001 S	.103 S	.001 T	.001 T	.001 S	.001 T	.002 F	.001 M	.001 W	.001 S	.087 M
15	.191 * T	.001 S	.006 S	.001 W	.001 F	.001 M	.001 W	.002 S	.001 T	.002 T	.001 S	.029 T
16	.084 F	.001 M	.001 M	.001 T	.001 S	.001 T	.001 T	.005 S	.001 W	.001 F	.001 M	.033 W
17	.015 S	.001 T	.001 T	.001 F	.001 S	.001 W	.001 F	.130 M	.001 T	.002 S	.001 T	.024 T
18	.001 S	.001 W	.001 W	.001 S	.001 M	.001 T	.001 S	.002 T	.001 F	.002 S	.001 W	.014 F
19	.001 M	.001 T	.002 T	.001 S	.001 T	.001 F	.001 S	.001 W	.001 S	.001 M	.001 T	.018 S
20	.016 T	.001 F	.001 F	.001 M	.001 W	.001 S	.001 M	.002 T	.001 S	.001 T	.001 F	.001 S
21	.042 W	.001 S	.021 S	.001 T	.001 T	.001 S	.001 T	.002 F	.002 M	.001 W	.001 S	.001 M
22	.014 T	.001 S	.001 S	.001 W	.001 F	.001 M	.001 W	.002 S	.001 T	.001 T	.001 S	.001 T
23	.053 F	.001 M	.001 M	.001 T	.001 S	.001 T	.001 T	.002 S	.002 W	.003 F	.001 M	.001 W
24	.018 S	.001 T	T	.001 F	.001 S	.001 W	.001 F	.001 M	.002 T	.001 S	.001 T	.001 T
25	.003 S	.001 W	.004 W	.001 S	.001 M	.001 T	.002 S	.001 T	.002 F	.001 S	.001 W	.001 F
26	.036 M	.001 T	.002 T	.012 S	.001 T	.001 F	.002 S	.002 W	.002 S	.001 M	.001 T	.001 S
27	.048 T	.001 F	.003 F	.076 M	.001 W	.001 S	.001 M	.001 T	.001 S	.001 T	.001 F	.070 S
28	.001 W	.019 S	.003 S	.001 T	.001 T	.001 S	.001 T	.002 F	.001 M	.001 W	.001 S	.036 M
29	.021 T		.003 S	.001 W	.001 F	.001 M	.001 W	.002 S	.001 T	.001 T	.001 S	.039 T
30	.003 F		.003 M	.001 T	.001 S	.001 T	.002 T	.002 S	.001 W	.001 F	.001 M	.053 W
31	.001 S		.003 T		.001 S		.001 F	.001 M		.001 S		.029 T
Number	22	28	30	30	31	30	31	31	30	31	30	31
Maximum	.258	.060	.103	.076	.001	.001	.002	.157	.002	.005	.302	.153
355 Total Samples 97.3 % Possible												
6 Starred (*) items exceeded the primary standard of .145 ppm												
Concentrations in parts per million (ppm)												

Frequency Distribution
Sulfur Dioxide Analyzer

Hawaii Volcanoes National Park

Monitoring Season: 01/01/98 - 12/31/98¹

Averaging Period	% Obs. ³	# Obs. ²	Min. Obs. ⁴	Percentile ⁵							Max. Obs.	2nd Max.	Arith. Mean	Geo. Mean	Geo. Stdv.
				10	30	50	70	90	95	99					
1-Hour	93	8129	0.001	0.001	0.001	0.001	0.001	0.005	0.037	0.281	0.952	0.949	0.0113	0.0017	3.27
3-Hour		2920	0.001	0.001	0.001	0.001	0.001	0.007	0.041	0.278	0.649	0.648			
24-Hour		365	0.001	0.001	0.001	0.001	0.002	0.031	0.057	0.150	0.290	0.237			

Concentrations in parts per million (ppm)

¹ Records for this report are selected in accordance with the AIRS Geo-Common file criteria. These criteria are based on the state-specific Monitoring Season defined in AIRS.


² The number of observations (# Obs.) includes all valid observations recorded within the Monitoring Season.

³ The percent of valid observations (% Obs.) is the percentage of valid days to the number of possible monitoring days during the Monitoring Season. A valid day is defined as a day with 9 or more valid observations between 9:00 a.m. and 9:00 p.m..


⁴ The minimum observation value (Min. Obs.) is the minimum daily maximum recorded during the Monitoring Season.

⁵ The percentiles and other statistics are derived from the daily maximums.


Second Highest 1-Hour Average Sulfur Dioxide Concentration




Highest 3-Hour Average Sulfur Dioxide Concentration




Highest 24-Hour Average Sulfur Dioxide Concentration




Legend




FIRST QUARTER (JAN-MAR)




SECOND QUARTER (APR-JUN)




THIRD QUARTER (JUL-SEP)




FOURTH QUARTER (OCT-DEC)



Maximum of Valid SO2 in ppb
 Indicates Data Exceeds Plot Scale
 Average of Valid SO2 in ppb
 (Bar is equal to +/- one standard deviation)
 Minimum of Valid SO2 in ppb




FIRST QUARTER (JAN-MAR)




94.0% Collected 34.8% Valid
2160 Possible /2031 Collected /751 Valid

SECOND QUARTER (APR-JUN)




95.0% Collected 85.1% Valid
2184 Possible /2074 Collected /1858 Valid

Sulfur Dioxide (ppb)




THIRD QUARTER (JUL-SEP)




95.2% Collected 95.2% Valid
2208 Possible /2102 Collected /2102 Valid

FOURTH QUARTER (OCT-DEC)



95.2% Collected 64.5% Valid
2208 Possible /2101 Collected /1425 Valid



95.6% Collected 64.2% Valid
8760 Possible /8373 Collected /5620 Valid

Sulfur Dioxide Precision Check Summary
Hawaii Volcanoes National Park

Precision checks are required by the Environmental Protection Agency (EPA) of all monitoring instruments collecting data which are to be submitted to the EPA Aerometric Information Retrieval System (AIRS). A precision check is performed by challenging the pollutant analyzer with a known concentration of gas (between 0.08 and 0.10 ppm for ozone and sulfur dioxide) from the pollutant transfer standard. This precision check must be performed at least every 14 days of monitoring operation. The percent difference between the analyzer and the transfer standard is then calculated. According to NPS Standard Operating Procedures, the pollutant analyzer must respond within 10% of the transfer standard. The table below gives the number of precision checks performed during each quarter, the average of all the individual precision check percent differences for the quarter, and the upper and lower 95% probability limits for precision checks. The probability limits represent the interval having a 95% chance of containing the true average percent difference. The quarterly average percent difference and probability limits should ideally be within +/- 10%.

Final Data 01/01/98 - 12/31/98				
Calendar Quarter	Number of Precision Checks	Average Percent Difference ^{1,2}	Lower 95% Probability Limit ³	Upper 95% Probability Limit ³
1	13	-7.86	-17.77	2.04
2	13	-1.03	-2.37	0.31
3	19	5.38	-1.26	12.02
4	13	3.85	-7.73	15.42

¹ Percent Difference = $\frac{\text{analyzer} - \text{transfer std}}{\text{transfer std}} \times 100$.

² Average Percent Difference is the mean of all individual precision check percent differences during the quarter.

³ Upper/Lower 95% Probability Limits = (Average Percent Difference) +/- (1.96)(Standard Deviation of precision check percent differences in the quarter.)

2.3 SUPPLEMENTAL SULFUR DIOXIDE DATA SUMMARY

Supplemental Sulfur Dioxide[‡]
 Five Highest Daily 1-Hour Averages, 3-Hour Block Averages,
 and 24-Hour Block Averages
 Hawaii Volcanoes National Park


Final Data
 01/01/98 - 12/31/98

Rank	Date	Hour	Concentration (ppb)
Highest Daily 1-Hour Averages			
1	11/08/98	15	1298
2	11/11/98	10	1063
3	08/11/98	8	978
4	11/12/98	9	875
5	08/17/98	12	835
Highest 3-Hour Block Averages			
1	11/11/98	9-11	766
2	11/11/98	6-8	603
3	11/05/98	9-11	537
4	08/11/98	9-11	529
5	11/08/98	15-17	528
Highest 24-Hour Block Averages			
1	11/11/98	00-23	346
2	11/08/98	00-23	188
3	08/11/98	00-23	169
4	12/08/98	00-23	157
5	11/12/98	00-23	145


[‡] Supplemental sulfur dioxide is collected from an additional analyzer.

Note: The primary and secondary ambient air standards for SO₂ are:
 Annual Primary Standard - 0.03 ppm; annual arithmetic mean not to be exceeded.
 (A value greater than .03 ppm, 34 ppb, or 80 ug/m³ exceeds the standard.)
 24-Hour Primary Standard - 0.14 ppm; not to be exceeded more than once per year.
 (A value greater than .14 ppm, 144 ppb, or 365 ug/m³ exceeds the standard.)
 3-Hour Secondary Standard - 0.5 ppm; not to be exceeded more than once per year.
 (A value greater than .5 ppm, 549 ppb, or 1300 ug/m³ exceeds the standard.)
 (40 CFR 50.4 and 50.5 with reference to Appendix A.)


Second Highest 1-Hour Average Supplemental Sulfur Dioxide Concentration




Highest 3-Hour Average Supplemental Sulfur Dioxide Concentration




Highest 24-Hour Average Supplemental Sulfur Dioxide Concentration




Legend




FIRST QUARTER (JAN-MAR)




SECOND QUARTER (APR-JUN)




THIRD QUARTER (JUL-SEP)




FOURTH QUARTER (OCT-DEC)



Maximum of Valid SO2ADD in ppb
 Average of Valid SO2ADD in ppb
 Minimum of Valid SO2ADD in ppb
 Indicates Data Exceeds Plot Scale
 (Bar is equal to +/- one standard deviation)




FIRST QUARTER (JAN-MAR)




97.0% Collected 10.9% Valid
2160 Possible /2096 Collected /235 Valid

SECOND QUARTER (APR-JUN)




95.0% Collected 85.1% Valid
2184 Possible /2074 Collected /1858 Valid

Supplemental Sulfur Dioxide (ppb)




THIRD QUARTER (JUL-SEP)




95.2% Collected 95.2% Valid
2208 Possible /2102 Collected /2102 Valid

FOURTH QUARTER (OCT-DEC)



95.2% Collected 64.5% Valid
2208 Possible /2101 Collected /1425 Valid



Supplemental Sulfur Dioxide Precision Check Summary †
Hawaii Volcanoes National Park

Precision checks are required by the Environmental Protection Agency (EPA) of all monitoring instruments collecting data which are to be submitted to the EPA Aerometric Information Retrieval System (AIRS). A precision check is performed by challenging the pollutant analyzer with a known concentration of gas (between 0.08 and 0.10 ppm for ozone and sulfur dioxide) from the pollutant transfer standard. This precision check must be performed at least every 14 days of monitoring¹ operation. The percent difference between the analyzer and the transfer standard is then calculated. According to NPS Standard Operating Procedures, the pollutant analyzer must respond within 10% of the transfer standard². The table below gives the number of precision checks performed during each quarter, the average³ of all the individual precision check percent differences for the quarter, and the upper and lower 95% probability limits for precision checks. The probability limits represent the interval having a 95% chance of containing the true average percent difference. The quarterly average percent difference and probability limits should ideally be within +/- 10%.

Final Data 01/01/98 - 12/31/98				
Calendar Quarter	Number of Precision Checks	Average Percent Difference ^{1,2}	Lower 95% Probability Limit ³	Upper 95% Probability Limit ³
1	173	-86.31	-148.28	-24.33
2	181	-85.16	-156.19	-14.12
3	184	-84.88	-158.01	-11.75
4	91	-81.33	-175.31	12.64

† Supplemental sulfur dioxide is collected from an additional analyzer.

¹ Percent Difference = $\frac{\text{analyzer} - \text{transfer std}}{\text{transfer std}} \times 100$.

² Average Percent Difference is the mean of all individual precision check percent differences during the quarter.

³ Upper/Lower 95% Probability Limits = (Average Percent Difference) +/- (1.96)(Standard Deviation of precision check percent differences in the quarter.)

2.4 METEOROLOGICAL DATA SUMMARY

Summary of Selected Meteorological Data

Hawaii Volcanoes National Park

Final Data

01/01/98 - 12/31/98


Parameter	Value	Units	Number	Std Dev
SCALAR WIND SPEED				
Average	4.2	m/s	7595	1.7
Maximum	10.2	m/s		
Percent calm = 2.04				
AMBIENT TEMPERATURE				
Average	14.9	degC	8591	2.5
Maximum	25.6	degC		
Minimum	7.2	degC		
RELATIVE HUMIDITY				
Average	93	percent	4611	9
Maximum	100	percent		
Minimum	53	percent		
PRECIPITATION (Rainfall or Snow melt)				
Average non-zero rate	1.0	mm/hr	2091	1.2
Maximum non-zero rate	18.8	mm/hr		
Minimum non-zero rate	.3	mm/hr		
Accumulated during period	2163.1	mm		
SOLAR RADIATION				
Average Daily Total	15,904,046	joules/m2day	365	4,691,650
Maximum Daily Total	25,817,600	joules/m2day		
Minimum Daily Total	2,659,200	joules/m2day		

Note: Calms are included in the average scalar wind speed and are defined as winds less than 0.5 m/s (1.0 mph).

Solar radiation terms are based on the calculation of the total amount of solar energy incident on a unit area during each day. The maximum and minimum daily totals are selected from the list of daily totals. The totals for all days are then added and divided by the number of days to yield the average daily total. Only days with 24 valid values are included in these statistics.


NA indicates instrument not available.

FIRST QUARTER (JAN-MAR)




99.0% Collected 46.1% Valid
2160 Possible /2139 Collected /995 Valid

SECOND QUARTER (APR-JUN)




99.8% Collected 89.5% Valid
2184 Possible /2180 Collected /1954 Valid

Scalar Wind Speed (m/s)




THIRD QUARTER (JUL-SEP)




100.0% Collected 100.0% Valid
2208 Possible /2208 Collected /2208 Valid

FOURTH QUARTER (OCT-DEC)



100.0% Collected 67.8% Valid
2208 Possible /2207 Collected /1497 Valid



99.7% Collected 76.0% Valid
8760 Possible /8734 Collected /6654 Valid

3.0 NATIONAL PARK SERVICE AIR RESOURCES DIVISION DATA SOURCES

3.1 GUIDE TO ATTACHED DATA DISKS

Data disks containing ASCII files of the validated hourly data, as shown in the following table are available. Please return the enclosed postcard or contact the address below. These data may be imported into other programs to perform additional data processing and analysis. The data format of each file is included within each file. The second table describes the validation codes used in the data tables to indicate why data are missing or invalid. Wind and pollutant frequency distribution tables in ASCII format are also included on the diskette if available for this site.

Data users should acknowledge the National Park Service Air Resources Division whenever using these data or any portion of this report.

3.2 OTHER SOURCES FOR RETRIEVING NATIONAL PARK SERVICE GASEOUS POLLUTANT DATA

The data contained in this report may also be obtained from the following sources:

- National Park Service AIRWeb (<http://www.aqd.nps.gov/natnet/ard>) - available after last quarter 1997
- EPA AIRS database
- Data requests directed to:

NPS Air Resources Division
Information Management Center
c/o Air Resource Specialists, Inc.
1901 Sharp Point Drive, Suite E
Fort Collins, Colorado 80525
Telephone: (970) 484-7941
Fax: (970) 484-3423
E-Mail: AIR-IMC@AIR-RESOURCE.COM

Data Disk Contents Summary	
File Name (s)	Description
Hourly	
ssssyy.DAT	All Validated Air Quality Data
ssssyymm.ppp	Monthly Data Summary Tables
ssssAN95.Rpp	Annual Wind and Pollutant Frequency Distribution
ssssQ195.Rpp	Quarter 1 Wind and Pollutant Frequency Distribution
ssssQ295.Rpp	Quarter 2 Wind and Pollutant Frequency Distribution
ssssQ395.Rpp	Quarter 3 Wind and Pollutant Frequency Distribution
ssssQ495.Rpp	Quarter 4 Wind and Pollutant Frequency Distribution
Where: ssss = site code yy = year mm = month ppp = air quality data parameter code AN = Annual Qn = Quarter 1-4 R = Wind Frequency distribution table	
CASTNet Weekly Species Summary Data	
File Name (s)	Description
CASTNet	
ssssCNyr.ASC	Weekly averages
Where: ssss = site code CN = CASTNet yr = year asc = ascii file	

NPS IMC and AIRS Invalid Data Codes			
NPS IMC VAL CODE	REASON	AIRS CODE	AIRS REASON
TO	Sample time out of limits	9973	Sample time out of limits
IW	Instrument warmup	9978	Voided by operator
OE	Operator error	9978	
BM	Begin monitoring	9979	Miscellaneous void
TL	Station temp low	9979	
OS	Off scale	9979	
EM	End monitoring	9979	
LI	Local interference	9979	
TH	Station temp high	9979	
IM	Instrument malfunction	9980	Machine malfunction
IN	Interference	9981	Bad weather
RF	Recording system failure	9983	Collection error
NA	No data	9987	Monitoring waived
PF	Power failure	9988	Power Failure
PC	Precision check	9990	Precision Check
ZS	Instrument zero/span check	9991	QC Control Points (Zero/Span)
SA	System audit	9992	QC Audit
PA	Performance audit	9992	
MT	Maintenance	9993	Maintenance/Routine Repairs
OR	Out for repair	9993	
CA	Calibration	9995	Multipoint calibration
SC	Station check	9998	Precision/zero/span

4.0 GLOSSARY

4.1 DEFINITIONS AND COMPUTATIONAL PROCEDURES FOR NATIONAL PARK SERVICE QUICK LOOK ANNUAL SUMMARY STATISTICS REPORT

The National Park Service Quick Look Annual Summary Statistics Table (Page 2-8) provides ozone summary statistics for various indices computed on a monthly basis for an entire year. Growing season (generically defined to be May 1 - September 30) and annual statistics are also presented under the "MAY-SEP" and "ANNUAL" columns, respectively. All concentrations are expressed in the units of parts per billion (PPB) and exposures in parts per billion-hours (PPB-HR). The definitions for each of the statistics appearing on the Quick Look Annual Summary Table are given below.


- (1) **Daily 1-Hr Maximum.** The maximum 1-hour average concentration recorded during each month, the growing season or the year regardless of the number of valid hourly observations recorded during a given day. The number in parentheses below this statistic, (N), indicates the number of days in the month, growing season, or year with valid data.
- (2) **Average Daily Maximum.** The average of all Daily 1-Hr Maxima during the month regardless of the number of Daily 1-Hr Maxima recorded during the month. For the "MAY-SEP" column the average of all the Daily Maxima recorded during the growing season is given. For the "ANNUAL" column the average of all the Daily Maxima is given. N is as in (1) above.
- (3) **Maximum Daily Mean.** The maximum of the valid daily means computed for each month, the growing season ("MAY-SEP" column), and the year ("ANNUAL" column). A valid daily mean is one for which 75% of the observations are available for each day, i.e., 18 hours. N is the number of days during each month, growing season, and year with at least 18 observations.
- (4) **Average Daily Mean.** The average of all valid daily means for the month, the growing season ("MAY-SEP" column), and the year ("ANNUAL" column). N is as in (3) above.
- (5) **Max Peak:Min Ratio.** The ratio of the Daily 1-Hr Maximum to the Daily 1-Hr Minimum. A ratio is computed only if a valid Daily Mean is computed and if the Daily 1-Hr Minimum is not equal to zero. N is the number of days with a valid Peak:Min ratio.
- (6) **Average Peak:Min Ratio.** The average of all Peak:Min ratios for the month, growing season, or year. N is as in (5) above.
- (7) **Max 9AM-4PM Average.** The maximum of all valid 9AM-4PM Averages computed for the month, growing season, or year. A valid 9AM-4PM Average is one which has 75% of the observations available during that time period (i.e., 6 hours. N is the number of days with valid averages.)

- (8) **Monthly 9AM-4PM Average.** The average of all valid 9AM-4PM Averages for the month, growing season, or year. N is as in (7) above.
- (9) **Max 7AM-7PM Average.** The maximum of all valid 7AM-7PM Averages computed for the month, growing season, or year. A valid 7AM-7PM Average is one which has 75% of the observations available during that time period, i.e., 9 hours. N is the number of days with valid averages.
- (10) **Monthly 7AM-7PM Average.** The average of all valid 7AM-7PM averages for the month, growing season, or year. N is as in (9) above.
- (11) **Monthly Mean.** The average of all 1-Hr ozone concentrations recorded during the month, growing season, or year. A mean is computed regardless of the number of hours with valid data. N is the number of hours with valid observations.
- (12) **SUM0 Exposure Index.** The monthly sum of all hourly ozone concentrations. Units are PPB-HR. The "MAY-SEP" column sums across the months of May through September to give the cumulative exposure for the growing season. The "ANNUAL" column sums across every month to give the cumulative exposure for the year. N is the number of hours with valid observations and is the same N as in (11) above.
- (13) **SUM60 Exposure Index.** The monthly sum of all hourly ozone concentrations equaling or exceeding 60 PPB. Units are PPB-HR. The "MAY-SEP" column sums across the months of May through September to give the cumulative exposure for the growing season. The "ANNUAL" column sums across every month to give the cumulative exposure for the year. N is the number of hours equaling or exceeding 60 PPB during the month, growing season, or year.
- (14) **SUM80 Exposure Index.** The monthly sum of all hourly ozone concentrations equaling or exceeding 80 PPB. Units are PPB-HR. The "MAY-SEP" column sums across the months of May through September to give the cumulative exposure for the growing season. The "ANNUAL" column sums across every month to give the cumulative exposure for the year. N is the number of hours equaling or exceeding 80 PPB during the month, growing season, or year.
- (15) **W126 Exposure Index.** The monthly sum of all hourly ozone concentrations where each concentration is weighted by a function that gives greater emphasis to the higher hourly concentrations while still including the lower ones. This weighting function provides a weighting value that is unique for each hourly ozone concentration. The weighting function, as described by Lefohn, Laurence, and Kohut¹ is:

$$w_i = \frac{1}{1 + 4403 \exp(-.126c_i)}$$

where

Weighting Function Used To Calculate W126 Exposure Index



w_i = weighting value for hourly concentration i ,
and
 c_i = hourly concentration i in PPB.

The graph of weighting value versus ozone concentration, in the figure to the left, illustrates the greater weights given to higher hourly ozone concentrations.

Each hour's weighting value is multiplied by its corresponding hourly concentration. This product is summed over all the valid hours in each month to calculate the monthly W126 exposure.

Thus, the monthly W126 exposure is:

$$W126 = \sum_{i=1}^n w_i c_i$$

where

W126 = monthly W126 exposure index,
 w_i = weighting value for hourly concentration i ,
 c_i = hourly concentration i in PPB, and
 n = number of hours in the month with valid ozone concentrations.

The "MAY-SEP" column sums across the months of May through September to give the cumulative exposure for the growing season. The "ANNUAL" column sums across every month to give the cumulative exposure for the year. The exposure units are PPB-HR.

Because each hour contributes to this exposure index, N is the number of hours with valid observations and is the same N as in (11) and (12) above.

The U.S. Environmental Protection Agency usually considers air quality statistics, such as a mean, to be "valid" (i.e., representative of the parameter being estimated for the time interval in question) only if 75% or more of the total possible observations have been measured during that time interval. Therefore, one should exercise caution when comparing these statistics between months and sites, particularly those that are not averages (e.g., maxima and exposures) whenever the number of valid observations is less than 75% of the total possible.

References

1. Lefohn, A.S., J. A. Laurence, and R. J. Kohut. 1988. A Comparison of Indices That Describe the Relationship Between Exposure to Ozone and Reduction in the Yield of Agricultural Crops. *Atmospheric Environment* 22, 1229-1240.

4.2 AIR QUALITY GLOSSARY

Acid Deposition: Air pollution produced when acid chemicals are incorporated into rain, snow, fog, or mist.

Aerometric Information Retrieval System (AIRS): A computer-based database of U.S. air pollution information administered by the EPA Office of Air Quality Planning and Standards (U.S. Environmental Protection Agency).

AIRWeb: Air Resources Web, an air quality information retrieval system for U.S. parks and wildlife refuges developed by the Air Resources Division of the National Park Service and the Air Quality Branch of the Fish and Wildlife Service.

Air Pollutant: An unwanted chemical or other material found in the air.

Air Pollution: Degradation of air quality resulting from unwanted chemicals or other materials occurring in the air.

Air Quality: The properties and degree of purity of air to which people and natural and heritage resources are exposed (in the context of national parks).

Air Pollution Control Permitting Process: Process by which facilities are permitted to emit specified types and quantities of air pollutants.

Air Quality Related Values (AQRVs): Values including visibility, flora, fauna, cultural and historical resources, odor, soil, water, and virtually all resources that are dependent upon and affected by air quality. "These values include visibility and those scenic, cultural, biological, and recreation resources of an area that are affected by air quality." (43 Fed. Reg. 15016)

Ambient Air: Air that is accessible to the public.

Class I: Areas of the country set aside under the Clean Air Act to receive the most stringent degree of air quality protection.

Class II: Areas of the country protected under the Clean Air Act but identified for somewhat less stringent protection from air pollution damage than Class I, except in specified cases.

Clean Air Act: Originally passed in 1963, our current national air pollution control program is based on the 1970 version of the law. Substantial revisions were made by the 1990 Clean Air Act Amendments.

Continuous Sampling Device: An air analyzer that measures air quality components continuously.

Criteria: Information on health and/or environmental effects of pollution (in the context of criteria air pollutants).

Criteria Air Pollutant: A group of very common air pollutants regulated by EPA on the basis of criteria and for which a National Ambient Air Quality Standard is established (SO₂, NO₂, PM₁₀, Pb, CO, O₃).

Emissions: Release of pollutants into the air from a source.

Environmental Protection Agency (EPA): The federal agency responsible for regulating air quality.

Monitoring: Measurement of air pollution.

National Ambient Air Quality Standards (NAAQS): Permissible levels of criteria air pollutant established to protect public health and welfare.

Ozone (O₃): A criteria air pollutant that is a strong oxidizing agent, reactive with many other compounds and surfaces, and a health hazard in high concentrations. Ozone is formed by nitrogen oxides and organic compounds reacting in sunlight.

Source: Any place or object from which air pollutants are released. Sources that are fixed in space are stationary sources; sources that move are mobile sources.

Sulfur Dioxide (SO₂): A criteria air pollutant that is a gas produced by burning coal and some industrial processes.

* Recent updates to this glossary may be found on the NPSARD AIRWeb - <http://www.aqd.nps.gov/natnet/ard/glossary.htm>.

4.3 GLOSSARY OF AIR QUALITY UNITS

Units Conversion Table			
Parameter Type	Multiply	By	To Obtain
Pollutant	ppm	1000	ppb
	ppm	1960	$\mu\text{g}/\text{m}^3$ Ozone (at 25°C)
	ppm	2615	$\mu\text{g}/\text{m}^3$ Sulfur Dioxide (at 25°C)
	ppb	0.001	ppm
	ppb	1.960	$\mu\text{g}/\text{m}^3$ Ozone (at 25°C)
	ppb	2.615	$\mu\text{g}/\text{m}^3$ Sulfur Dioxide (at 25°C)
	$\mu\text{g}/\text{m}^3$ Ozone (25°C)	0.0005102	ppm
	$\mu\text{g}/\text{m}^3$ Ozone (25°C)	0.5102	ppb
	$\mu\text{g}/\text{m}^3$ Sulfur Dioxide (25°C)	0.0003824	ppm
	$\mu\text{g}/\text{m}^3$ Sulfur Dioxide (25°C)	0.3824	ppb
Wind Speed	m/s	2.05	mph
	mph	0.489	m/s
Solar Radiation	ly/min	697	w/m^2
	w/m^2	0.00143	ly/min
Precipitation	mm/hr	0.0394	in/hr
	in/hr	25.4	mm/hr
Temperature	$^{\circ}\text{C} + 17.78$	1.8	$^{\circ}\text{F}$
	$^{\circ}\text{F} - 32$	5/9	$^{\circ}\text{C}$
<p>Where:</p> <p>ppm = parts per million</p> <p>ppb = parts per billion</p> <p>$\mu\text{g}/\text{m}^3$ = micrograms per cubic meter (at 25°C)</p> <p>m/s = meters per second</p> <p>mps = miles per hour</p> <p>ly/min = langley's per minute</p> <p>w/m^2 = watts per square meter</p> <p>mm/hr = millimeters per hour</p> <p>in/hr = inches per hour</p> <p>$^{\circ}\text{C}$ = degrees centigrade</p> <p>$^{\circ}\text{F}$ = degrees fahrenheit</p>			