

Annual Data Summary
DEATH VALLEY NATIONAL PARK
1998
National Park Service
Gaseous Air Pollutant Monitoring Network



AIR RESOURCES DIVISION
RESEARCH AND MONITORING BRANCH
12795 West Alameda Parkway
P.O. Box 25287
Lakewood, Colorado 80225
Telephone: (303) 969-2820
Fax: (303) 969-2822

This Annual Data Summary was prepared under NPS Contract CX-1270-96-007 by:

Air Resource Specialists, Inc.
1901 Sharp Point Drive, Suite E
Fort Collins, Colorado 80525
Telephone: (970) 484-7941
Fax: (970) 484-3423

For additional copies of this report or reports for other NPS units, contact:

National Park Service Air Resources Division
Information Management Center
c/o Air Resource Specialists, Inc.
1909 Sharp Point Drive, Suite E
Fort Collins, Colorado 80525
Telephone: (970) 484-7941
Fax: (970) 484-3423
E-Mail: AIR-IMC@AIR-RESOURCE.COM

or

National Park Service
Air Resources Division
P.O. Box 25287
Lakewood, Colorado 80225-02587
Telephone: (303) 969-2130
E-Mail: AQ_INFO@AQD.NPS.GOV

ACKNOWLEDGEMENTS

The National Park Service Air Resources Division (ARD) recognizes the level of effort required by individual park units, site operators, auditors, cooperating state and local agencies, and ARD contractors. ARD sincerely appreciates the contributions of all participants in assisting with the collection, validation, and reporting of these air quality and meteorological data.

At Death Valley National Park, the ARD specifically recognizes Dick Anderson and Arnie Peterson for performing the technical and administrative skills required to help produce the data presented within this report.

TABLE OF CONTENTS

| <u>Section</u> | | <u>Page</u> |
|-----------------------|---|--------------------|
| 1.0 | INTRODUCTION | 1-1 |
| 1.1 | The National Park Service Gaseous Pollutant Monitoring Network | 1-1 |
| 1.2 | Death Valley National Park | 1-3 |
| 2.0 | DATA SUMMARY | 2-1 |
| 2.1 | Overview | 2-1 |
| 2.2 | Ozone Data Summary | 2-7 |
| 2.3 | Meteorological Data Summary | 2-23 |
| 2.4 | Dry Deposition Data Summary | 2-27 |
| 3.0 | NATIONAL PARK SERVICE AIR RESOURCES DIVISION DATA SOURCES | 3-1 |
| 3.1 | Guide to Attached Data Disks | 3-1 |
| 3.2 | Other Sources for Retrieving National Park Service Gaseous Pollutant Data | 3-1 |
| 4.0 | GLOSSARY | 4-1 |
| 4.1 | Definitions and Computational Procedures for National Park Service Quick Look Annual Summary Statistics Table | 4-1 |
| 4.2 | Air Quality Glossary | 4-4 |

LIST OF FIGURES AND TABLES

| | <u>Page</u> |
|--|--------------------|
| Map of National Park Service Gaseous Pollutant Monitoring Network Ozone and Sulfur Dioxide Monitoring Sites | 1-2 |
| Site Specifications | 1-4 |
| Data Collection Statistics | 2-2 |
| First Quarter Data Stackplot | 2-3 |
| Second Quarter Data Stackplot | 2-4 |
| Third Quarter Data Stackplot | 2-5 |
| Fourth Quarter Data Stackplot | 2-6 |

LIST OF FIGURES AND TABLES (Continued)

| | <u>Page</u> |
|--|-------------|
| Ozone Quick Look Annual Summary Statistics | 2-8 |
| Ozone Annual Frequency Distribution | 2-9 |
| Ozone Daily 1-Hour Maximum Concentrations and National Ambient Air Quality Standards Comparison | 2-10 |
| Attainment Status with EPA Proposed Primary Ozone Standards | 2-11 |
| Ten Highest 1-Hour Average Ozone Concentrations | 2-12 |
| Episodes with 1-Hour Ozone Concentrations 100ppb and > 124 ppb | 2-13 |
| Episodes with 8-Hour Ozone Concentrations >84 ppb | 2-14 |
| Ozone Rank Listings of Second Highest 1-Hour Average Concentrations, Maximum 8-Hour Average Concentrations, and Annual Sum60 Exposure Index for All NPS Monitoring Sites | 2-15 |
| Plot of Maximum Ozone Concentration Comparison for Three Years | 2-16 |
| Map of National Park Service Gaseous Pollutant Monitoring Network, Second Highest Hourly Ozone Concentration | 2-17 |
| Quarterly Diurnal Ozone Plots | 2-18 |
| Annual Diurnal Ozone Plot | 2-19 |
| Quarterly Ozone Pollutant Roses | 2-20 |
| Annual Ozone Pollutant Roses | 2-21 |
| Ozone Precision Check Data Summary | 2-22 |
| Summary of Selected Meteorological Data | 2-24 |
| Quarterly Wind Roses | 2-25 |
| Annual Wind Roses | 2-26 |
| Quarterly and Annual Average Concentrations | 2-28 |
| Weekly Concentrations Report | 2-29 |
| Three Year Comparison of Maximum and Average Concentrations | 2-30 |

LIST OF FIGURES AND TABLES (Continued)

| | <u>Page</u> |
|--|--------------------|
| Average Particulate Nitrate Concentrations | 2-31 |
| Nitric Acid Concentrations | 2-32 |
| Average Total Nitrate Concentrations | 2-33 |
| Average Ammonium Concentrations | 2-34 |
| Average Particulate Sulfate Concentrations | 2-35 |
| Average Sulfur Dioxide Concentrations | 2-36 |
| SO ₄ /SO ₂ Ratio | 2-37 |
| Data Disk Contents | 3-2 |
| NPS IMC and AIRS Invalid Data Codes | 3-3 |

1.0 INTRODUCTION

1.1 THE NATIONAL PARK SERVICE GASEOUS POLLUTANT MONITORING NETWORK

Gaseous air pollutants, including ozone and sulfur dioxide, are of concern to the National Park Service (NPS). Pollutants like these can affect park unit biological resources as well as the health of park unit residents and visitors. The NPS established a gaseous pollutant monitoring program for several pollutants linked to effects on NPS resources. This program was designed to meet certain resource management objectives.

The primary objective of this monitoring program is to establish the status and trends of park unit air quality conditions and to determine if a park unit is exceeding the National Ambient Air Quality Standards established by the U.S. Environmental Protection Agency (EPA) to protect public health and welfare. In addition, such monitoring is designed to detect changes or trends in pollution levels over time. A monitoring station may also be established if there is documented biological injury due to air pollution in a park unit. Information on ambient air pollution levels is an important part of research on effects of air pollutants on NPS resources, and can help confirm suspected causes of observed effects.

Other monitoring objectives call for the collection of data to support the National Park Service's required involvement in both the development of state air quality control plans, and the evaluation of permit applications for new or expanding air pollution sources wishing to locate near park units. The Clean Air Act gives federal land managers and superintendents an affirmative responsibility to protect air quality related values in Class I areas and to assess whether new sources will have an adverse impact on park unit resources and values. Information on air quality levels in NPS units can also be used to evaluate the performance of atmospheric models that simulate how pollutants are transported into park units and predict impacts on the park unit caused by air pollution sources.

The National Park Service Gaseous Pollutant Monitoring Network site locations and measured parameters collected in this reporting year are shown on the map on the following page. During this reporting period, 40 monitoring sites in 35 units of the National Park System had some combination of ozone, sulfur dioxide, meteorological, and CASTNet dry deposition monitoring. Monitoring methods and quality assurance procedures used in the national park network meet the applicable 40 CFR Part 58 EPA requirements. This allows for the direct comparison of NPS collected data with that collected by the EPA, and state and local air pollution control agencies. Data collected by this network are incorporated in the EPA Aerometric Information Retrieval System (AIRS) database which is a national database of all air quality data collected throughout the country. These data are also stored in the NPS Air Resources Division's Information Management Center (IMC) that allows for easy access and analysis of data.

This report includes a variety of data summaries for data collected at an individual monitoring site at a national park unit during this reporting period. These summaries highlight the average range and frequency of the data collected during the year. A PC-compatible diskette containing a digital copy of all data collected during the year and data summary products included in this report is available. Individual reports are generated for each site where monitoring was conducted in the national park network.

**NATIONAL PARK SERVICE
GASEOUS POLLUTANT MONITORING NETWORK**

1998 Ozone and Sulfur Dioxide Monitoring Sites






1.2 DEATH VALLEY NATIONAL PARK

Death Valley National Park is located along the California and Nevada state line near the western edge of the Basin and Range Province. Its location and site specifications are presented on the following page.

Death Valley National Park was proclaimed by Congress on February 11, 1933. The park was designated a Biosphere reserve in 1984.

The park, over 2,000,000 acres in size, is a large desert nearly surrounded by high mountains and contains the lowest point in the western hemisphere. In 1994 the park was increased to 3.3 million acres.

**Death Valley
National Park
Ozone Monitoring
NPS Gaseous Air Pollutant Monitoring Network**



| SITE IDENTIFICATION | | MAP INFORMATION | |
|-------------------------|-------------------|-----------------|---------------|
| Site Abbreviation: | DEVA | Mean Elevation: | 125 m |
| AIRS ID NO.: | 06-027-0101 | Longitude: | 116° 50' 52"W |
| INSTRUMENTATION | | Latitude: | 36° 30' 31"N |
| | | UTM Zone: | 11 |
| O ₃ Analyzer | Delta Temperature | Easting: | 513631 m |
| Calibrator | Temperature | Northing: | 4040172 m |
| Wind Speed | Solar Radiation | Map Reference: | Death Valley |
| Wind Direction | Precipitation | | NJ 11-11 |
| Relative Humidity | | | 1,250,000 |

2.0 DATA SUMMARY

2.1 OVERVIEW

Based on the site specifications during this annual reporting period, data summaries and statistics are provided in this section.

Data Collection Statistics
Death Valley National Park

Final Data

01/01/98 - 12/31/98

| Parameter | Par Code | Data Recovery | | | Valid Data | |
|---------------------------------------|----------|---------------|---------------|-------------|------------|---------|
| | | No. Possible | No. Collected | % Collected | No. Valid | % Valid |
| Ozone Analyzer | O3 | 8760 | 8138 | 92.9 | 7947 | 90.7 |
| Scalar Wind Speed | SWS | 8760 | 8608 | 98.3 | 8481 | 96.8 |
| Vector Wind Speed | VWS | 8760 | 8607 | 98.3 | 8480 | 96.8 |
| Vector Wind Direction | VWD | 8760 | 8607 | 98.3 | 8480 | 96.8 |
| Standard Deviation for Wind Direction | SDWD | 8760 | 8596 | 98.1 | 8469 | 96.7 |
| Ambient Temperature (aspirated) | TMP | 8760 | 8607 | 98.3 | 8607 | 98.3 |
| Delta Temperature | DTP | 8760 | 8607 | 98.3 | 8607 | 98.3 |
| Relative Humidity | RH | 8760 | 8633 | 98.6 | 3783 | 43.2 |
| Precipitation | RNF | 8760 | 8571 | 97.8 | 8571 | 97.8 |
| Wetness Sensor | WET | 8760 | 8610 | 98.3 | 8610 | 98.3 |
| Solar Radiation | SOL | 8760 | 8614 | 98.3 | 8614 | 98.3 |
| Filter Pack Flow Rate | FLOW | 8760 | 8582 | 98.0 | 8582 | 98.0 |

Notes: All statistics are for hourly averages.

The number collected does not include normal maintenance or events beyond the control of the network.

The percent valid is calculated against the number possible.

Automatic zeros and spans are performed daily on most ambient gas analyzers, therefore, no ambient data can be collected during this time. As a result, the maximum percent valid for ambient gas data typically can not be greater than 95.8.

NPS Performance Goals:


Quarterly Criteria:

100% of sites, >= 85% valid data capture
90% of sites, >= 90% valid data capture
80% of sites, >= 95% valid data capture

Monthly Criteria:

100% of sites, >= 60% valid data capture
90% of sites, >= 75% valid data capture
80% of sites, >= 85% valid data capture

Death Valley National Park




Final Validation

First Quarter 1998

deva-pv.stk - deva98.dat 08-20-1999

Death Valley National Park




Final Validation

Second Quarter 1998

deva-pv.stk - deva98.dat 08-20-1999

Death Valley National Park




Final Validation

Third Quarter 1998

deva-pv.stk - deva98.dat 08-20-1999

Death Valley National Park



Final Validation

Fourth Quarter 1998

deva-pv.stk - deva98.dat 08-20-1999

2.2 OZONE DATA SUMMARY

Ozone Quick Look Annual Summary Statistics
Death Valley National Park

01/01/98 - 12/31/98

| STATISTIC | JAN | FEB | MAR | APR | MAY | JUN | JUL | AUG | SEP | OCT | NOV | DEC | MAY-SEP | ANNUAL |
|-------------------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|---------|--------|
| DAILY 1-HR MAXIMUM | 50 | 50 | 68 | 82 | 74 | 85 | 92 | 87 | 78 | 79 | 53 | 50 | 92 | 92 |
| NO. OF DAYS | (31) | (28) | (31) | (30) | (31) | (30) | (31) | (29) | (29) | (31) | (30) | (31) | (150) | (362) |
| AVERAGE DAILY MAXIMUM | 39 | 44 | 51 | 57 | 62 | 66 | 74 | 65 | 64 | 55 | 44 | 41 | 66 | 55 |
| NO. OF DAYS | (31) | (28) | (31) | (30) | (31) | (30) | (31) | (29) | (29) | (31) | (30) | (31) | (150) | (362) |
| MAXIMUM DAILY MEAN | 43 | 45 | 56 | 67 | 67 | 75 | 81 | 65 | 68 | 67 | 45 | 45 | 81 | 81 |
| NO. OF DAYS | (30) | (28) | (31) | (30) | (30) | (30) | (24) | (24) | (25) | (30) | (30) | (31) | (133) | (343) |
| AVERAGE DAILY MEAN | 36 | 40 | 44 | 51 | 53 | 58 | 65 | 56 | 56 | 49 | 40 | 37 | 57 | 48 |
| NO. OF DAYS | (30) | (28) | (31) | (30) | (30) | (30) | (24) | (24) | (25) | (30) | (30) | (31) | (133) | (343) |
| MAX PEAK:MIN RATIO | 1.429 | 1.515 | 1.625 | 1.771 | 2.226 | 1.973 | 1.673 | 1.740 | 1.622 | 1.639 | 1.514 | 1.704 | 2.226 | 2.226 |
| NO. OF DAYS | (30) | (28) | (31) | (30) | (30) | (30) | (24) | (24) | (25) | (30) | (30) | (31) | (133) | (343) |
| AVERAGE PEAK:MIN RATIO | 1.238 | 1.284 | 1.341 | 1.337 | 1.425 | 1.395 | 1.334 | 1.423 | 1.399 | 1.285 | 1.276 | 1.230 | 1.396 | 1.328 |
| NO. OF DAYS | (30) | (28) | (31) | (30) | (30) | (30) | (24) | (24) | (25) | (30) | (30) | (31) | (133) | (343) |
| MAX 9AM-4PM AVERAGE | 43 | 47 | 63 | 75 | 72 | 82 | 84 | 74 | 72 | 73 | 51 | 46 | 84 | 84 |
| NO. OF DAYS | (29) | (28) | (28) | (29) | (30) | (29) | (24) | (26) | (27) | (30) | (27) | (29) | (136) | (336) |
| MONTHLY 9AM-4PM AVERAGE | 36 | 41 | 46 | 54 | 57 | 61 | 67 | 58 | 60 | 52 | 41 | 37 | 60 | 51 |
| NO. OF DAYS | (29) | (28) | (28) | (29) | (30) | (29) | (24) | (26) | (27) | (30) | (27) | (29) | (136) | (336) |
| MAX 7AM-7PM AVERAGE | 44 | 46 | 60 | 73 | 71 | 80 | 83 | 71 | 70 | 70 | 49 | 47 | 83 | 83 |
| NO. OF DAYS | (30) | (28) | (31) | (30) | (30) | (30) | (23) | (23) | (27) | (30) | (30) | (31) | (133) | (343) |
| MONTHLY 7AM-7PM AVERAGE | 36 | 40 | 46 | 53 | 56 | 60 | 67 | 57 | 58 | 51 | 41 | 38 | 59 | 50 |
| NO. OF DAYS | (30) | (28) | (31) | (30) | (30) | (30) | (23) | (23) | (27) | (30) | (30) | (31) | (133) | (343) |
| MONTHLY MEAN | 36 | 40 | 44 | 51 | 53 | 58 | 65 | 56 | 56 | 49 | 40 | 37 | 57 | 48 |
| NO. OF HOURS | (691) | (630) | (695) | (677) | (691) | (675) | (611) | (588) | (619) | (697) | (676) | (697) | (3184) | (7947) |
| SUM0 EXPOSURE INDEX | 24573 | 24910 | 30725 | 34389 | 36948 | 38905 | 39603 | 32782 | 34361 | 34331 | 27008 | 26024 | 182599 | 384559 |
| NO. OF HOURS | (691) | (630) | (695) | (677) | (691) | (675) | (611) | (588) | (619) | (697) | (676) | (697) | (3184) | (7947) |
| SUM60 EXPOSURE INDEX | - | - | 1075 | 3430 | 12135 | 16962 | 29324 | 11967 | 13013 | 5912 | - | - | 83401 | 93818 |
| NO. OF HOURS | (0) | (0) | (17) | (51) | (190) | (248) | (422) | (183) | (200) | (92) | (0) | (0) | (1243) | (1403) |
| SUM80 EXPOSURE INDEX | - | - | - | 82 | - | 1970 | 4333 | 250 | - | - | - | - | 6553 | 6635 |
| NO. OF HOURS | (0) | (0) | (0) | (1) | (0) | (24) | (52) | (3) | (0) | (0) | (0) | (0) | (79) | (80) |
| W126 EXPOSURE INDEX | 608 | 935 | 2532 | 5602 | 8601 | 12822 | 19220 | 8784 | 9102 | 5541 | 1118 | 844 | 58529 | 75710 |
| NO. OF HOURS | (691) | (630) | (695) | (677) | (691) | (675) | (611) | (588) | (619) | (697) | (676) | (697) | (3184) | (7947) |

Concentrations in parts per billion (ppb)

* Statistics defined in the Quick Look subsection of the Glossary

Exposures in parts per billion-hours (ppb-hr)

Frequency Distribution

Ozone Analyzer

Death Valley National Park

Monitoring Season: 01/01/98 - 12/31/98¹

| Averaging Period | % Obs. ³ | # Obs. ² | Min. Obs. ⁴ | Percentile ⁵ | | | | | | | Max. Obs. | 2nd Max. | Arith. Mean | Geo. Mean | Geo. Stdv. |
|---|---------------------|---------------------|---------------------------|-------------------------|-------|-------|-------|-------|-------|-------|--------------|-------------|----------------|--------------|---------------|
| | | | | 10 | 30 | 50 | 70 | 90 | 95 | 99 | | | | | |
| 1-Hour | 93 | 7947 | 0.030 | 0.040 | 0.046 | 0.053 | 0.061 | 0.073 | 0.079 | 0.085 | 0.092 | 0.087 | 0.0547 | 0.0532 | 1.26 |
| Concentrations in parts per million (ppm) | | | | | | | | | | | | | | | |

¹ Records for this report are selected in accordance with the AIRS Geo-Common file criteria. These criteria are based on the state-specific Monitoring Season defined in AIRS.

² The number of observations (# Obs.) includes all valid observations recorded within the Monitoring Season.

³ The percent of valid observations (% Obs.) is the percentage of valid days to the number of possible monitoring days during the Monitoring Season. A valid day is defined as a day with 9 or more valid observations between 9:00 a.m. and 9:00 p.m..

⁴ The minimum observation value (Min. Obs.) is the minimum daily maximum recorded during the Monitoring Season.

⁵ The percentiles and other statistics are derived from the daily maximums.

Ozone Standards Report and
Daily Maximum 1-Hour Concentrations (ppm)
Death Valley National Park

01/01/98 - 12/31/98

| Day | Jan-98 | Feb-98 | Mar-98 | Apr-98 | May-98 | Jun-98 | Jul-98 | Aug-98 | Sep-98 | Oct-98 | Nov-98 | Dec-98 |
|------------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| 1 | .034 T | .044 S | .042 S | .057 W | .067 F | .071 M | .069 W | .069 S | .058 T | .057 T | .048 S | .050 T |
| 2 | .045 F | .043 M | .046 M | .048 T | .058 S | .068 T | .085 T | .057 S | .061 W | .066 F | .048 M | .044 W |
| 3 | .044 S | .046 T | .056 T | .059 F | .061 S | .060 W | .082 F | .050 M | .063 T | .060 S | .046 T | .042 T |
| 4 | .044 S | .041 W | .047 W | .054 S | .053 M | .053 T | .087 S | .049 T | .058 F | .049 S | .042 W | .049 F |
| 5 | .044 M | .042 T | .058 T | .053 S | .052 T | .056 F | .080 S | W | .059 S | .047 M | .050 T | .043 S |
| 6 | .042 T | .045 F | .050 F | .051 M | .054 W | .084 S | .077 M | T | .062 S | .046 T | .045 F | .045 S |
| 7 | .039 W | .044 S | .046 S | .051 T | .057 T | .062 S | .065 T | F | .058 M | .059 W | .051 S | .044 M |
| 8 | .035 T | .043 S | .048 S | .054 W | .055 F | .056 M | W | S | .062 T | .079 T | .047 S | .044 T |
| 9 | .034 F | .042 M | .050 M | .056 T | .061 S | .056 T | T | .065 S | .053 W | .070 F | .045 M | .047 W |
| 10 | .030 S | .040 T | .047 T | .067 F | .062 S | .060 W | .076 F | .073 M | T | .061 S | .040 T | .046 T |
| 11 | .034 S | .044 W | .044 W | .062 S | .072 M | .068 T | .071 S | .055 T | .072 F | .065 S | .043 W | .041 F |
| 12 | M | .040 T | .052 T | .053 S | .055 T | .068 F | .072 S | W | .073 S | .067 M | .042 T | .037 S |
| 13 | T | .044 F | .057 F | .056 M | .054 W | .064 S | .081 M | .079 T | .061 S | .065 T | .042 F | .035 S |
| 14 | .033 W | .044 S | .052 S | .053 T | .054 T | .057 S | .076 T | .087 F | .054 M | .063 W | .038 S | .045 M |
| 15 | .037 T | .048 S | .047 S | .056 W | .065 F | .062 M | W | .072 S | T | .055 T | .037 S | .043 T |
| 16 | .038 F | .045 M | .053 M | .055 T | S | .064 T | T | .075 S | .062 W | .048 F | .038 M | .040 W |
| 17 | .036 S | .043 T | .050 T | .058 F | .060 S | .051 W | F | .060 M | .070 T | .045 S | .053 T | .036 T |
| 18 | .036 S | .047 W | .051 W | .055 S | .067 M | .073 T | S | .074 T | .073 F | .048 S | .043 W | .035 F |
| 19 | .044 M | .040 T | .050 T | .057 S | .068 T | .083 F | S | .063 W | .065 S | .049 M | .042 T | .045 S |
| 20 | .041 T | .048 F | .052 F | .060 M | .069 W | .080 S | .080 M | .069 T | .066 S | T | .040 F | .045 S |
| 21 | .044 W | .047 S | .068 S | .055 T | .060 T | .068 S | .083 T | .060 F | .070 M | .056 W | .039 S | .040 M |
| 22 | .040 T | .046 S | .061 S | .073 W | .058 F | .064 M | .073 W | S | .078 T | .059 T | .039 S | .039 T |
| 23 | .038 F | .040 M | .049 M | .057 T | .065 S | .082 T | .092 T | .061 S | .075 W | .059 F | .041 M | .042 W |
| 24 | .036 S | .050 T | .060 T | .053 F | .069 S | .085 W | .079 F | .070 M | .076 T | .055 S | .047 T | .039 T |
| 25 | .035 S | .047 W | .043 W | .052 S | .061 M | .071 T | .065 S | .070 T | .068 F | .049 S | .048 W | .038 F |
| 26 | .035 M | .046 T | .043 T | .056 S | .056 T | .065 F | .059 S | W | .066 S | .046 M | .047 T | .034 S |
| 27 | .039 T | .044 F | .045 F | .059 M | .061 W | .075 S | .056 M | .069 T | .051 S | .042 T | .044 F | .031 S |
| 28 | .041 W | .044 S | .050 S | .060 T | .059 T | .069 S | .064 T | .060 F | .056 M | .053 W | .051 S | .031 M |
| 29 | .050 T | | .048 S | .062 W | .069 F | .054 M | W | S | .068 T | .052 T | .042 S | .032 T |
| 30 | .043 F | | .047 M | .082 T | .066 S | .062 T | .075 T | S | .063 W | .043 F | .049 M | .032 W |
| 31 | .039 S | | .062 T | | .074 S | | .062 F | M | | .048 S | | .046 T |
| Valid Days | 29 | 28 | 31 | 30 | 30 | 30 | 23 | 21 | 28 | 30 | 30 | 31 |
| Maximum | .050 | .050 | .068 | .082 | .074 | .085 | .092 | .087 | .078 | .079 | .053 | .050 |
| Violations | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

| | |
|------------------------|---|
| 7947 Total Samples | 0 Daily-maxima exceeding the standard of .12 ppm (starred[*]) |
| 90.7 % Possible | 8 Missing days assumed to be less than the standard |
| 341 Valid daily maxima | 0 Daily maximas exceed the alert level of .200 ppm |

Concentrations in parts per million (ppm)

Death Valley National Park

1998 Attainment Status With U.S. Environmental Protection Agency (EPA) PRIMARY Ozone National Ambient Air Quality Standard

Ozone Season: January through December

The primary National Ambient Air Quality Standard for ozone is designed to protect human health. The level of the primary ozone standard promulgated by the EPA on July 18, 1997 is 0.08 parts per million (ppm) [80 parts per billion, (ppb)], daily maximum 8-hour average. The primary ozone standard is met at an ambient monitoring site when the 3-year average of the annual fourth-highest daily maximum 8-hour average ozone concentration is less than or equal to 0.08 ppm. This standard is not met when the 3-year average is greater than 0.08 ppm. Using the EPA's rounding convention, a computed 3-year average ozone concentration of 0.085 ppm (85 ppb) is the smallest value that is greater than the level of the 0.08 ppm standard.

The primary standard requires 90 percent data completeness, on average, during the 3-year period, with no single year within the period having less than 75 percent data completeness. This data completeness requirement would have to be satisfied in order to determine that the standard has been met at a monitoring site. However, calendar years with less than 75 percent data completeness are included in the computation if the annual fourth-highest daily maximum 8-hour concentration is greater than the level of the standard. A site could be found not to have met the standard with less than complete data. The percent data completeness is the percent of valid ozone monitoring days. A day is valid if valid 8-hour averages are available for at least 75 percent of possible hours in the day (i.e., at least 18 of the 24 averages). An 8-hour average is considered valid if at least 75 percent (or 6) of the hourly averages for the 8-hour period are available.

The table below lists the 3-year average fourth-highest daily maximum 8-hour ozone concentration based on data collected during the reported year and the two previous years. This is the number to compare to the level of the new primary standard. The 3-year average data completeness percent and the reported year highest five daily maximum 8-hour averages are also tabulated. A 'No' in the Data Comp % Met? column indicates EPA data completeness requirement was not met for the three-year period.

| Year | 3-Year Avg 4th High Daily Max 8-hr Ozone (ppb) | 3-Year Avg Data Complete % | Data Complete % Met? | Annual 1st High Daily Max 8-hr Ozone (ppb) | Annual 2nd High Daily Max 8-hr Ozone (ppb) | Annual 3rd High Daily Max 8-hr Ozone (ppb) | Annual 4th High Daily Max 8-hr Ozone (ppb) | Annual 5th High Daily Max 8-hr Ozone (ppb) |
|------|--|----------------------------|----------------------|--|--|--|--|--|
| 1998 | 79 | 90% | Yes | 85 | 84 | 82 | 82 | 82 |

Ozone
 Ten Highest Daily 1-Hour Average Maximum Concentrations
 Death Valley National Park

Final Data
 01/01/98 - 12/31/98

| Rank | Date | Hour | Concentration (ppb) |
|------|----------|------|------------------------|
| 1 | 07/23/98 | 16 | 92* |
| 2 | 07/19/98 | 11 | 88* |
| 3 | 07/04/98 | 18 | 87* |
| 4 | 08/14/98 | 16 | 87 |
| 5 | 06/24/98 | 14 | 85* |
| 6 | 07/02/98 | 12 | 85* |
| 7 | 06/06/98 | 18 | 84* |
| 8 | 07/17/98 | 12 | 84* |
| 9 | 06/19/98 | 11 | 83 |
| 10 | 07/21/98 | 16 | 83* ** |

* Other high value(s) were also recorded during one or more hours in the day.

** This value was also recorded on one or more days later in the reporting period.

Episodes with 1-Hour Ozone Concentrations
 > 100 ppb and > 124 ppb
 Death Valley National Park

Final Data
 01/01/98 - 12/31/98

| Date | Beginning Hour | No. Hours | | Max (ppb) |
|---|----------------|-----------|----------|-----------|
| | | > 100 ppb | >124 ppb | |
| No values exceeded 100 ppb during this period | | | | |
| Total | | 0 | 0 | |

Note: The primary and secondary national ambient air standard for ozone that applied in 1996 is 0.12 ppm over a one hour period not to be exceeded more than once per year. (A value greater than .12 ppm, 124 ppb, or 235 ug/m³ exceeds the standard.) (40 CFR 50.9 with reference to Appendix D and H.)

**Episodes with 8-Hour Average Ozone Concentrations > 84 ppb
Death Valley National Park**

**Final Data
01/01/98 - 12/31/98**

| Date | Start and End Time of Daily Maximum 8-Hour Average > 84 ppb (hr) | Daily Maximum 8-Hour Average (ppb) | Number of 8-Hour Averages > 84 ppb During the Day |
|--|---|--|---|
| No values exceeded 84 ppb during this period | | | |
| 0 | Days with 8-hour average concentrations > 84 ppb | | |

Note: This table presents episodes of high ozone based on running 8-hour averages. In 1997, the EPA published new primary and secondary national ambient air quality standards for ozone based on 8-hour average ozone concentrations. Attainment of the new primary standard is reached if the annual fourth highest daily maximum 8-hour ozone concentration, averaged over three years, does not exceed 0.08 ppm (84 ppb or 157 ug/m). (40 CFR 50.10.)





Ozone Rank Listings of Second Highest 1-Hour Average Concentrations, 4th Highest 8-Hour Average Concentrations, and Annual SUM60 Exposure Index for All NPS Monitoring Sites

01/01/98 - 12/31/98


| Second Highest 1-Hour Average Concentration | | |
|---|------|---------------------|
| Site | Rank | Concentration (ppb) |
| JOTR-YV | 1 | 138 |
| GRSM-LR | 2 | 134 |
| SEKI-LK | 3 | 125 |
| SHEN-BM | 4 | 124 |
| ACAD-CM | 5 | 123 |
| GRSM-CM | 6 | 123 |
| CHAM-XX | 7 | 122 |
| COWP-XX | 8 | 122 |
| GRSM-CD | 9 | 117 |
| SEKI-LP | 10 | 117 |
| CACO-XX | 11 | 115 |
| MACA-HM | 12 | 114 |
| PINN-ES | 13 | 113 |
| COSW-XX | 14 | 106 |
| GRSM-CC | 15 | 105 |
| YOSE-TD | 16 | 105 |
| ROMO-LP | 17 | 100 |
| SAGU-PC | 18 | 94 |
| LAVO-ML | 19 | 91 |
| EVER-BC | 20 | 90 |
| DEVA-PV | 21 | 88 |
| GRBA-MY | 22 | 83 |
| CHIS-XX | 23 | 81 |
| CHIR-ES | 24 | 80 |
| BIBE-KB | 25 | 78 |
| CANY-IS | 26 | 78 |
| VOYA-SB | 27 | 78 |
| GRCA-AS | 28 | 76 |
| MEVE-MY | 29 | 73 |
| YELL-WT | 30 | 72 |
| CRMO-VC | 31 | 69 |
| MORA-TW | 32 | 69 |
| GLAC-WG | 33 | 63 |
| OLYM-VC | 34 | 62 |
| THRO-VC | 35 | 60 |
| DENA-HQ | 36 | 57 |
| NOCA-MM | 37 | 53 |
| VIIS-LP | 38 | 49 |

| 4th Highest 8-hour Average Concentration | | |
|--|------|---------------------|
| Site | Rank | Concentration (ppb) |
| GRSM-LR | 1 | 110 |
| JOTR-YV | 2 | 110 |
| GRSM-CM | 3 | 107 |
| SHEN-BM | 4 | 107 |
| GRSM-CD | 5 | 106 |
| SEKI-LP | 6 | 99 |
| MACA-HM | 7 | 98 |
| COWP-XX | 8 | 97 |
| ACAD-CM | 9 | 95 |
| SEKI-LK | 10 | 94 |
| YOSE-TD | 11 | 94 |
| CHAM-XX | 12 | 89 |
| PINN-ES | 13 | 88 |
| GRSM-CC | 14 | 86 |
| CACO-XX | 15 | 84 |
| COSW-XX | 16 | 82 |
| DEVA-PV | 17 | 82 |
| ROMO-LP | 18 | 80 |
| LAVO-ML | 19 | 78 |
| SAGU-PC | 20 | 77 |
| GRCA-AS | 21 | 73 |
| CANY-IS | 22 | 72 |
| EVER-BC | 23 | 72 |
| GRBA-MY | 24 | 71 |
| BIBE-KB | 25 | 70 |
| CHIR-ES | 26 | 68 |
| MEVE-MY | 27 | 68 |
| VOYA-SB | 28 | 68 |
| YELL-WT | 29 | 67 |
| CHIS-XX | 30 | 66 |
| CRMO-VC | 31 | 66 |
| GLAC-WG | 32 | 58 |
| DENA-HQ | 33 | 55 |
| THRO-VC | 34 | 55 |
| MORA-TW | 35 | 51 |
| OLYM-VC | 36 | 46 |
| NOCA-MM | 37 | 43 |
| VIIS-LP | 38 | 39 |


| Annual Sum60 Exposure Index | | | |
|-----------------------------|------|-------------|------|
| Site | Rank | Sum60 Count | |
| GRSM-CM | 1 | 198342 | 2702 |
| GRSM-CD | 2 | 187437 | 2577 |
| SHEN-BM | 3 | 170745 | 2387 |
| GRSM-LR | 4 | 164447 | 2231 |
| JOTR-YV | 5 | 127317 | 1769 |
| DEVA-PV | 6 | 93818 | 1403 |
| YOSE-TD | 7 | 92922 | 1338 |
| SEKI-LP | 8 | 92214 | 1230 |
| SEKI-LK | 9 | 84666 | 1144 |
| MACA-HM | 10 | 82293 | 1162 |
| COWP-XX | 11 | 70877 | 970 |
| CANY-IS | 12 | 68500 | 1075 |
| GRCA-AS | 13 | 63994 | 996 |
| ROMO-LP | 14 | 59083 | 897 |
| SAGU-PC | 15 | 57929 | 869 |
| GRSM-CC | 16 | 52679 | 742 |
| ACAD-CM | 17 | 45061 | 638 |
| CACO-XX | 18 | 44769 | 651 |
| PINN-ES | 19 | 43209 | 609 |
| CHIR-ES | 20 | 35885 | 565 |
| GRBA-MY | 21 | 35229 | 551 |
| LAVO-ML | 22 | 33289 | 501 |
| MEVE-MY | 23 | 32220 | 511 |
| CHAM-XX | 24 | 31595 | 434 |
| BIBE-KB | 25 | 26226 | 409 |
| COSW-XX | 26 | 26019 | 364 |
| CRMO-VC | 27 | 17194 | 274 |
| EVER-BC | 28 | 16065 | 239 |
| YELL-WT | 29 | 9932 | 157 |
| CHIS-XX | 30 | 9696 | 150 |
| VOYA-SB | 31 | 8985 | 137 |
| GLAC-WG | 32 | 1407 | 23 |
| MORA-TW | 33 | 638 | 10 |
| OLYM-VC | 34 | 307 | 5 |
| THRO-VC | 35 | 181 | 3 |
| DENA-HQ | 36 | 0 | 0 |
| NOCA-MM | 37 | 0 | 0 |
| VIIS-LP | 38 | 0 | 0 |




FIRST QUARTER (JAN-MAR)




SECOND QUARTER (APR-JUN)




THIRD QUARTER (JUL-SEP)



FOURTH QUARTER (OCT-DEC)



Maximum of Valid O3 in ppb
 Average of Valid O3 in ppb (Bar is equal to +/- one standard deviation)
 Minimum of Valid O3 in ppb
 Indicates Data Exceeds Plot Scale




Death Valley
National Park

Quarterly Ozone
Pollutant Rose


1998

FIRST QUARTER (JAN-MAR)




93.2% Collected 93.1% Valid
2160 Possible /2014 Collected /2012 Valid

SECOND QUARTER (APR-JUN)




93.4% Collected 93.4% Valid
2184 Possible /2040 Collected /2040 Valid

THIRD QUARTER (JUL-SEP)




88.6% Collected 77.3% Valid
2208 Possible /1957 Collected /1706 Valid

FOURTH QUARTER (OCT-DEC)



93.7% Collected 91.7% Valid
2208 Possible /2068 Collected /2024 Valid



92.2% Collected 88.8% Valid
8760 Possible /8079 Collected /7782 Valid

Ozone Precision Check Summary Death Valley National Park

Precision checks are required by the Environmental Protection Agency (EPA) of all monitoring instruments collecting data which are to be submitted to the EPA Aerometric Information Retrieval System (AIRS). A precision check is performed by challenging the pollutant analyzer with a known concentration of gas (between 0.08 and 0.10 ppm for ozone and sulfur dioxide) from the pollutant transfer standard. This precision check must be performed at least every 14 days of monitoring operation. The percent difference between the analyzer and the transfer standard is then calculated. According to NPS Standard Operating Procedures, the pollutant analyzer must respond within 10% of the transfer standard.² The table below gives the number of precision checks performed during each quarter, the average of all the individual precision check percent differences for the quarter, and the upper and lower 95% probability limits for precision checks. The probability limits represent the interval having a 95% chance of containing the true average percent difference. The quarterly average percent difference and probability limits should ideally be within +/- 10%.

| Final Data | | | | |
|---------------------|----------------------------|---|--|--|
| 01/01/98 - 12/31/98 | | | | |
| Calendar Quarter | Number of Precision Checks | Average Percent Difference ^{1,2} | Lower 95% Probability Limit ³ | Upper 95% Probability Limit ³ |
| 1 | 15 | -5.89 | -19.52 | 7.75 |
| 2 | 13 | -2.38 | -6.25 | 1.50 |
| 3 | 13 | -0.71 | -6.10 | 4.68 |
| 4 | 13 | -3.42 | -7.14 | 0.29 |

¹ Percent Difference = $\frac{\text{analyzer} - \text{transfer std}}{\text{transfer std}} \times 100$.

² Average Percent Difference is the mean of all individual precision check percent differences during the quarter.

³ Upper/Lower 95% Probability Limits = (Average Percent Difference) +/- (1.96)(Standard Deviation of precision check percent differences in the quarter.)

2.3 METEOROLOGICAL DATA SUMMARY

Summary of Selected Meteorological Data

Death Valley National Park

Final Data

01/01/98 - 12/31/98


| Parameter | Value | Units | Number | Std Dev |
|--|------------|--------------|--------|-----------|
| SCALAR WIND SPEED | | | | |
| Average | 3.4 | m/s | 8481 | 2.7 |
| Maximum | 16.8 | m/s | | |
| Percent calm = 0.13 | | | | |
| AMBIENT TEMPERATURE | | | | |
| Average | 24.1 | degC | 8607 | 10.0 |
| Maximum | 50.1 | degC | | |
| Minimum | 3.5 | degC | | |
| RELATIVE HUMIDITY | | | | |
| Average | 26 | percent | 3783 | 12 |
| Maximum | 88 | percent | | |
| Minimum | 5 | percent | | |
| PRECIPITATION (Rainfall or Snow melt) | | | | |
| Average non-zero rate | 1.1 | mm/hr | 94 | 1.0 |
| Maximum non-zero rate | 5.1 | mm/hr | | |
| Minimum non-zero rate | .3 | mm/hr | | |
| Accumulated during period | 107.4 | mm | | |
| SOLAR RADIATION | | | | |
| Average Daily Total | 17,235,357 | joules/m2day | 365 | 6,623,347 |
| Maximum Daily Total | 27,734,400 | joules/m2day | | |
| Minimum Daily Total | 1,686,400 | joules/m2day | | |

Note: Calms are included in the average scalar wind speed and are defined as winds less than 0.5 m/s (1.0 mph).

Solar radiation terms are based on the calculation of the total amount of solar energy incident on a unit area during each day. The maximum and minimum daily totals are selected from the list of daily totals. The totals for all days are then added and divided by the number of days to yield the average daily total. Only days with 24 valid values are included in these statistics.


NA indicates instrument not available.

FIRST QUARTER (JAN-MAR)




99.6% Collected 99.6% Valid
2160 Possible /2151 Collected /2151 Valid

SECOND QUARTER (APR-JUN)




99.4% Collected 99.4% Valid
2184 Possible /2171 Collected /2171 Valid

Scalar Wind Speed (m/s)




THIRD QUARTER (JUL-SEP)




94.2% Collected 90.6% Valid
2208 Possible /2079 Collected /2000 Valid

FOURTH QUARTER (OCT-DEC)



99.9% Collected 97.7% Valid
2208 Possible /2206 Collected /2158 Valid



98.3% Collected 96.8% Valid
8760 Possible /8607 Collected /8480 Valid

2.4 DRY DEPOSITION DATA SUMMARY

Clean Air Status and Trends Network (CASTNet) Dry Deposition Monitoring

In 1995, the National Park Service (NPS) and the Environmental Protection Agency (EPA) entered a partnership to jointly measure dry deposition in park units, mostly in the West. A portion of the 1997 data collected from this partnership is presented in this section.

Atmospheric deposition of acidic species takes two pathways: wet deposition and dry deposition. Wet deposition is the result of precipitation events (rain, snow, or fog) that remove particles and gases from the atmosphere. Dry deposition is less event driven, but still involves the transfer of particles and gases from the atmosphere to surfaces and plants. Wet deposition has been well documented for many years. In the national parks, the National Acidic Deposition Program (NADP) measures and reports wet deposition (see the web site at <http://nadp.sws.uiuc.edu> for further information). Dry deposition is much harder to measure and a smaller network of monitoring stations is involved. The method used to measure dry deposition is sometimes called the "inferential method" because air quality concentration data are combined with meteorological measurements and land use functions to compute deposition velocities. The CASTNet program provides long-term estimates of total acidic deposition by adding dry deposition values to wet deposition values.

This annual summary report presents the air quality concentration portion of the dry deposition inferential method, which is the only currently available data set. These data were compiled from the analyses of filters collected by CASTNet deposition filter pack systems in the parks. The filter pack analyses yielded weekly average concentrations of particulate sulfate (SO_4^{2-}), particulate nitrate (NO_3^-), particulate ammonium (NH_4^+), sulfur dioxide (SO_2), and nitric acid (HNO_3). In some cases, the positive ions Na^+ , K^+ , Ca^{2+} , and Mg^{2+} were also measured from the filter samples. These concentration data for the individual ionic species are presented as weekly bar charts and summarized by quarter and by year in this report. Concentration data can be used to compare sites and to indicate the amount of acidic species available for deposition. As with the continuous analyzer data, the filter pack concentration data are included on a computer diskette that accompanies this report.

Estimated dry deposition values derived from EPA modeling will be reported at a later time to complete the inferential analyses. When available, these modeling results will be posted on the NPS Air Resources Division Internet web site at <http://www.aqd.nps.gov/ard1> or on the EPA CASTNet site (<http://www.epa.gov/ardpublic/acidrain/castnet/about.html>). Initial CASTNet results have shown that dry deposition can be a significant portion of total acidic deposition.

CASTNet Dry Deposition Monitoring
Quarterly and Annual Average Concentrations
Death Valley National Park
1/1/98 - 12/31/98

| Quarter | No. Valid Samples | p-NO ₃ (ug/m ³) | HNO ₃ (ug/m ³) | Total NO ₃ (ug/m ³) | NH ₄ (ug/m ³) | p-SO ₄ (ug/m ³) | SO ₂ (ug/m ³) | SO ₄ /SO ₂ Ratio |
|--------------------|-------------------|---|--|---|---|---|---|---|
| 1 | 13 | 0.295 | 1.029 | 1.308 | 0.150 | 0.587 | 0.177 | 3.324 |
| 2 | 13 | 0.388 | 1.450 | 1.815 | 0.320 | 0.977 | 0.466 | 2.098 |
| 3 | 13 | 0.436 | 2.641 | 3.036 | 0.540 | 1.604 | 0.680 | 2.359 |
| 4 | 13 | 0.228 | 1.233 | 1.442 | 0.199 | 0.605 | 0.261 | 2.323 |
| Annual Average | | 0.337 | 1.588 | 1.900 | 0.302 | 0.943 | 0.396 | 2.384 |
| Standard Deviation | | 0.184 | 0.888 | 0.989 | 0.194 | 0.582 | 0.279 | |

| Data Recovery Table | | | |
|---------------------|-----------------|--------------|-----------------|
| Total No. Filters | No. Invalidated | Data Capture | No. Valid Hours |
| 52 | 0 | 100.0% | 8545.0 |


CASTNet Dry Deposition Monitoring Weekly Concentrations Report
 Death Valley National Park
 1/1/98 - 12/31/98

| On Date | Off Date | p-NO ₃ (ug/m ³) | HNO ₃ (ug/m ³) | Total NO ₃ (ug/m ³) | NH ₄ (ug/m ³) | p-SO ₄ (ug/m ³) | SO ₂ (ug/m ³) | SO ₄ /SO ₂ Ratio |
|----------|----------|---|--|---|---|---|---|---|
| 12/30/97 | 01/06/98 | 0.251 | 0.969 | 1.205 | 0.118 | 0.729 | 0.170 | 4.286 |
| 01/06/98 | 01/13/98 | 0.488 | 1.725 | 2.186 | 0.213 | 0.319 | 0.150 | 2.122 |
| 01/13/98 | 01/20/98 | 0.227 | 1.150 | 1.359 | 0.108 | 0.362 | 0.147 | 2.461 |
| 01/20/98 | 01/27/98 | 0.105 | 0.441 | 0.539 | 0.067 | 0.172 | 0.129 | 1.328 |
| 01/27/98 | 02/03/98 | 0.433 | 1.515 | 1.924 | 0.141 | 0.400 | 0.230 | 1.744 |
| 02/03/98 | 02/10/98 | 0.505 | 0.449 | 0.947 | 0.074 | 0.222 | 0.243 | 0.912 |
| 02/10/98 | 02/17/98 | 0.338 | 0.758 | 1.084 | 0.130 | 0.408 | 0.187 | 2.183 |
| 02/17/98 | 02/24/98 | 0.213 | 0.693 | 0.895 | 0.075 | 0.342 | 0.171 | 1.995 |
| 02/24/98 | 03/03/98 | 0.179 | 0.777 | 0.944 | 0.127 | 0.308 | 0.145 | 2.128 |
| 03/03/98 | 03/10/98 | 0.374 | 1.038 | 1.395 | 0.184 | 2.125 | 0.159 | 13.393 |
| 03/10/98 | 03/17/98 | 0.144 | 1.564 | 1.683 | 0.364 | 1.008 | 0.239 | 4.222 |
| 03/17/98 | 03/24/98 | 0.231 | 1.546 | 1.752 | 0.219 | 0.648 | 0.182 | 3.555 |
| 03/24/98 | 03/31/98 | 0.351 | 0.756 | 1.095 | 0.130 | 0.594 | 0.145 | 4.092 |
| 03/31/98 | 04/07/98 | 0.520 | 1.330 | 1.828 | 0.191 | 1.346 | 0.314 | 4.286 |
| 04/07/98 | 04/14/98 | 0.366 | 1.548 | 1.890 | 0.241 | 0.905 | 0.381 | 2.375 |
| 04/14/98 | 04/21/98 | 0.139 | 0.467 | 0.599 | 0.182 | 0.531 | 0.182 | 2.913 |
| 04/21/98 | 04/28/98 | 0.115 | 0.252 | 0.363 | 0.096 | 0.266 | 0.409 | 0.650 |
| 04/28/98 | 05/05/98 | 0.504 | 1.450 | 1.931 | 0.273 | 0.993 | 0.201 | 4.950 |
| 05/05/98 | 05/12/98 | 0.431 | 1.348 | 1.757 | 0.346 | 1.003 | 0.242 | 4.148 |
| 05/12/98 | 05/19/98 | 0.357 | 1.366 | 1.702 | 0.255 | 0.780 | 0.433 | 1.802 |
| 05/19/98 | 05/26/98 | 0.356 | 1.523 | 1.855 | 0.381 | 1.043 | 0.588 | 1.775 |
| 05/26/98 | 06/02/98 | 0.472 | 2.010 | 2.450 | 0.288 | 0.824 | 0.560 | 1.471 |
| 06/02/98 | 06/09/98 | 0.359 | 1.655 | 1.988 | 0.366 | 1.085 | 0.454 | 2.391 |
| 06/09/98 | 06/16/98 | 0.232 | 1.669 | 1.875 | 0.372 | 0.985 | 0.336 | 2.932 |
| 06/16/98 | 06/23/98 | 0.765 | 1.670 | 2.408 | 0.467 | 1.273 | 0.832 | 1.530 |
| 06/23/98 | 06/30/98 | 0.428 | 2.555 | 2.942 | 0.702 | 1.664 | 1.121 | 1.484 |
| 06/30/98 | 07/07/98 | 0.492 | 2.512 | 2.965 | 0.427 | 1.354 | 0.972 | 1.393 |
| 07/07/98 | 07/14/98 | 0.345 | 2.515 | 2.821 | 0.637 | 1.636 | 1.069 | 1.530 |
| 07/14/98 | 07/21/98 | 0.585 | 2.383 | 2.930 | 0.656 | 1.930 | 1.164 | 1.659 |
| 07/21/98 | 07/28/98 | 0.466 | 1.967 | 2.402 | 0.627 | 1.954 | 0.650 | 3.008 |
| 07/28/98 | 08/04/98 | 0.261 | 1.841 | 2.073 | 0.278 | 0.791 | 0.532 | 1.488 |
| 08/04/98 | 08/11/98 | 0.306 | 2.470 | 2.737 | 0.522 | 1.454 | 0.732 | 1.985 |
| 08/11/98 | 08/18/98 | 0.323 | 2.635 | 2.916 | 0.469 | 1.401 | 0.664 | 2.108 |
| 08/18/98 | 08/25/98 | 0.360 | 2.468 | 2.788 | 0.364 | 1.057 | 0.553 | 1.909 |
| 08/25/98 | 09/01/98 | 0.473 | 4.290 | 4.695 | 0.887 | 2.733 | 0.791 | 3.455 |
| 09/01/98 | 09/08/98 | 0.160 | 1.906 | 2.036 | 0.582 | 1.827 | 0.232 | 7.862 |
| 09/08/98 | 09/15/98 | 0.220 | 2.588 | 2.766 | 0.425 | 1.223 | 0.263 | 4.655 |
| 09/15/98 | 09/22/98 | 0.621 | 2.708 | 3.286 | 0.541 | 1.602 | 0.720 | 2.226 |
| 09/22/98 | 09/29/98 | 1.060 | 4.056 | 5.051 | 0.605 | 1.890 | 0.497 | 3.804 |
| 09/29/98 | 10/06/98 | 0.226 | 1.756 | 1.954 | 0.334 | 0.931 | 0.239 | 3.902 |
| 10/06/98 | 10/13/98 | 0.330 | 3.198 | 3.477 | 0.316 | 0.985 | 0.540 | 1.822 |
| 10/13/98 | 10/20/98 | 0.248 | 1.585 | 1.807 | 0.261 | 0.899 | 0.347 | 2.595 |
| 10/20/98 | 10/27/98 | 0.224 | 1.553 | 1.752 | 0.332 | 0.894 | 0.425 | 2.105 |
| 10/27/98 | 11/03/98 | 0.102 | 0.914 | 1.001 | 0.186 | 0.503 | 0.197 | 2.547 |
| 11/03/98 | 11/10/98 | 0.188 | 0.822 | 0.997 | 0.156 | 0.389 | 0.387 | 1.004 |
| 11/10/98 | 11/17/98 | 0.097 | 1.135 | 1.213 | 0.138 | 0.297 | 0.221 | 1.340 |
| 11/17/98 | 11/24/98 | 0.109 | 0.871 | 0.966 | 0.222 | 0.610 | 0.204 | 2.995 |
| 11/24/98 | 12/01/98 | 0.526 | 1.734 | 2.232 | 0.253 | 0.765 | 0.280 | 2.733 |
| 12/01/98 | 12/08/98 | 0.236 | 1.107 | 1.325 | 0.078 | 0.315 | 0.179 | 1.757 |
| 12/08/98 | 12/15/98 | 0.168 | 0.297 | 0.460 | 0.063 | 0.248 | 0.109 | 2.270 |
| 12/15/98 | 12/22/98 | 0.391 | 0.611 | 0.992 | 0.137 | 0.686 | 0.155 | 4.434 |
| 12/22/98 | 12/29/98 | 0.125 | 0.444 | 0.563 | 0.112 | 0.347 | 0.105 | 3.301 |


Death Valley National Park

CASTNet Dry Deposition Monitoring
 Three Year Comparison of Maximum and Average Concentrations


p-NO₃




HNO₃




Total NO₃




NH₄




p-SO₄



SO₂








Key:

| | |
|-------------|--------------------|
| ACAD | Acadia NP |
| BIBE | Big Bend NP |
| CANY | Canyonlands NP |
| CHIR | Chiricahua NM |
| DENA | Denali NP |
| DEVA | Death Valley NP |
| EVER | Everglades NP |
| GLAC | Glacier NP |
| GRBA | Great Basin NP |
| GRCA | Grand Canyon NP |
| GRSM | Great Smokies NP |
| JOTR | Joshua Tree NP |
| LAVO | Lassen Volcanic NP |
| MEVE | Mesa Verde NP |
| MORA | Mount Rainier NP |
| NOCA | North Cascades NP |
| OLYM | Olympic NP |
| PINN | Pinnacles NM |
| ROMO | Rocky Mountain NP |
| SEKI | Sequoia NP |
| SHEN | Shenandoah NP |
| THRO | Th. Roosevelt NP |
| VIIS | Virgin Islands NP |
| VOYA | Voyageurs NP |
| YELL | Yellowstone NP |
| YOSE | Yosemite NP |




Key:

| | |
|-------------|--------------------|
| ACAD | Acadia NP |
| BIBE | Big Bend NP |
| CANY | Canyonlands NP |
| CHIR | Chiricahua NM |
| DENA | Denali NP |
| DEVA | Death Valley NP |
| EVER | Everglades NP |
| GLAC | Glacier NP |
| GRBA | Great Basin NP |
| GRCA | Grand Canyon NP |
| GRSM | Great Smokies NP |
| JOTR | Joshua Tree NP |
| LAVO | Lassen Volcanic NP |
| MEVE | Mesa Verde NP |
| MORA | Mount Rainier NP |
| NOCA | North Cascades NP |
| OLYM | Olympic NP |
| PINN | Pinnacles NM |
| ROMO | Rocky Mountain NP |
| SEKI | Sequoia NP |
| SHEN | Shenandoah NP |
| THRO | Th. Roosevelt NP |
| VIIS | Virgin Islands NP |
| VOYA | Voyageurs NP |
| YELL | Yellowstone NP |
| YOSE | Yosemite NP |


CASTNet Dry Deposition Monitoring Sites / National Park Service Units

Average Ammonium (NH₄) Concentrations (ug/m³) for Entire Year 1998




Key:

| | |
|-------------|--------------------|
| ACAD | Acadia NP |
| BIBE | Big Bend NP |
| CANY | Canyonlands NP |
| CHIR | Chiricahua NM |
| DENA | Denali NP |
| DEVA | Death Valley NP |
| EVER | Everglades NP |
| GLAC | Glacier NP |
| GRBA | Great Basin NP |
| GRCA | Grand Canyon NP |
| GRSM | Great Smokies NP |
| JOTR | Joshua Tree NP |
| LAVO | Lassen Volcanic NP |
| MEVE | Mesa Verde NP |
| MORA | Mount Rainier NP |
| NOCA | North Cascades NP |
| OLYM | Olympic NP |
| PINN | Pinnacles NM |
| ROMO | Rocky Mountain NP |
| SEKI | Sequoia NP |
| SHEN | Shenandoah NP |
| THRO | Th. Roosevelt NP |
| VIIS | Virgin Islands NP |
| VOYA | Voyageurs NP |
| YELL | Yellowstone NP |
| YOSE | Yosemite NP |




Key:

| | |
|-------------|--------------------|
| ACAD | Acadia NP |
| BIBE | Big Bend NP |
| CANY | Canyonlands NP |
| CHIR | Chiricahua NM |
| DENA | Denali NP |
| DEVA | Death Valley NP |
| EVER | Everglades NP |
| GLAC | Glacier NP |
| GRBA | Great Basin NP |
| GRCA | Grand Canyon NP |
| GRSM | Great Smokies NP |
| JOTR | Joshua Tree NP |
| LAVO | Lassen Volcanic NP |
| MEVE | Mesa Verde NP |
| MORA | Mount Rainier NP |
| NOCA | North Cascades NP |
| OLYM | Olympic NP |
| PINN | Pinnacles NM |
| ROMO | Rocky Mountain NP |
| SEKI | Sequoia NP |
| SHEN | Shenandoah NP |
| THRO | Th. Roosevelt NP |
| VIIS | Virgin Islands NP |
| VOYA | Voyagers NP |
| YELL | Yellowstone NP |
| YOSE | Yosemite NP |



Key:

| | |
|-------------|--------------------|
| ACAD | Acadia NP |
| BIBE | Big Bend NP |
| CANY | Canyonlands NP |
| CHIR | Chiricahua NM |
| DENA | Denali NP |
| DEVA | Death Valley NP |
| EVER | Everglades NP |
| GLAC | Glacier NP |
| GRBA | Great Basin NP |
| GRCA | Grand Canyon NP |
| GRSM | Great Smokies NP |
| JOTR | Joshua Tree NP |
| LAVO | Lassen Volcanic NP |
| MEVE | Mesa Verde NP |
| MORA | Mount Rainier NP |
| NOCA | North Cascades NP |
| OLYM | Olympic NP |
| PINN | Pinnacles NM |
| ROMO | Rocky Mountain NP |
| SEKI | Sequoia NP |
| SHEN | Shenandoah NP |
| THRO | Th. Roosevelt NP |
| VIIS | Virgin Islands NP |
| VOYA | Voyageurs NP |
| YELL | Yellowstone NP |
| YOSE | Yosemite NP |



Key:

| | |
|-------------|--------------------|
| ACAD | Acadia NP |
| BIBE | Big Bend NP |
| CANY | Canyonlands NP |
| CHIR | Chiricahua NM |
| DENA | Denali NP |
| DEVA | Death Valley NP |
| EVER | Everglades NP |
| GLAC | Glacier NP |
| GRBA | Great Basin NP |
| GRCA | Grand Canyon NP |
| GRSM | Great Smokies NP |
| JOTR | Joshua Tree NP |
| LAVO | Lassen Volcanic NP |
| MEVE | Mesa Verde NP |
| MORA | Mount Rainier NP |
| NOCA | North Cascades NP |
| OLYM | Olympic NP |
| PINN | Pinnacles NM |
| ROMO | Rocky Mountain NP |
| SEKI | Sequoia NP |
| SHEN | Shenandoah NP |
| THRO | Th. Roosevelt NP |
| VIIS | Virgin Islands NP |
| VOYA | Voyageurs NP |
| YELL | Yellowstone NP |
| YOSE | Yosemite NP |

3.0 NATIONAL PARK SERVICE AIR RESOURCES DIVISION DATA SOURCES

3.1 GUIDE TO ATTACHED DATA DISKS

Data disks containing ASCII files of the validated hourly data, as shown in the following table are available. Please return the enclosed postcard or contact the address below. These data may be imported into other programs to perform additional data processing and analysis. The data format of each file is included within each file. The second table describes the validation codes used in the data tables to indicate why data are missing or invalid. Wind and pollutant frequency distribution tables in ASCII format are also included on the diskette if available for this site.

Data users should acknowledge the National Park Service Air Resources Division whenever using these data or any portion of this report.

3.2 OTHER SOURCES FOR RETRIEVING NATIONAL PARK SERVICE GASEOUS POLLUTANT DATA

The data contained in this report may also be obtained from the following sources:

- National Park Service AIRWeb (<http://www.aqd.nps.gov/natnet/ard>) - available after last quarter 1997
- EPA AIRS database
- Data requests directed to:

NPS Air Resources Division
Information Management Center
c/o Air Resource Specialists, Inc.
1901 Sharp Point Drive, Suite E
Fort Collins, Colorado 80525
Telephone: (970) 484-7941
Fax: (970) 484-3423
E-Mail: AIR-IMC@AIR-RESOURCE.COM

| Data Disk Contents Summary | |
|--|---|
| File Name (s) | Description |
| Hourly | |
| ssssyy.DAT | All Validated Air Quality Data |
| ssssyymm.ppp | Monthly Data Summary Tables |
| ssssAN95.Rpp | Annual Wind and Pollutant Frequency Distribution |
| ssssQ195.Rpp | Quarter 1 Wind and Pollutant Frequency Distribution |
| ssssQ295.Rpp | Quarter 2 Wind and Pollutant Frequency Distribution |
| ssssQ395.Rpp | Quarter 3 Wind and Pollutant Frequency Distribution |
| ssssQ495.Rpp | Quarter 4 Wind and Pollutant Frequency Distribution |
| Where: ssss = site code yy = year mm = month ppp = air quality data parameter code AN = Annual Qn = Quarter 1-4 R = Wind Frequency distribution table | |
| CASTNet Weekly Species Summary Data | |
| File Name (s) | Description |
| CASTNet | |
| ssssCNyr.ASC | Weekly averages |
| Where: ssss = site code CN = CASTNet yr = year asc = ascii file | |

| NPS IMC and AIRS Invalid Data Codes | | | |
|--|----------------------------|------------------|-------------------------------|
| NPS IMC VAL CODE | REASON | AIRS CODE | AIRS REASON |
| TO | Sample time out of limits | 9973 | Sample time out of limits |
| IW | Instrument warmup | 9978 | Voided by operator |
| OE | Operator error | 9978 | |
| BM | Begin monitoring | 9979 | Miscellaneous void |
| TL | Station temp low | 9979 | |
| OS | Off scale | 9979 | |
| EM | End monitoring | 9979 | |
| LI | Local interference | 9979 | |
| TH | Station temp high | 9979 | |
| IM | Instrument malfunction | 9980 | Machine malfunction |
| IN | Interference | 9981 | Bad weather |
| RF | Recording system failure | 9983 | Collection error |
| NA | No data | 9987 | Monitoring waived |
| PF | Power failure | 9988 | Power Failure |
| PC | Precision check | 9990 | Precision Check |
| ZS | Instrument zero/span check | 9991 | QC Control Points (Zero/Span) |
| SA | System audit | 9992 | QC Audit |
| PA | Performance audit | 9992 | |
| MT | Maintenance | 9993 | Maintenance/Routine Repairs |
| OR | Out for repair | 9993 | |
| CA | Calibration | 9995 | Multipoint calibration |
| SC | Station check | 9998 | Precision/zero/span |

4.0 GLOSSARY

4.1 DEFINITIONS AND COMPUTATIONAL PROCEDURES FOR NATIONAL PARK SERVICE QUICK LOOK ANNUAL SUMMARY STATISTICS REPORT

The National Park Service Quick Look Annual Summary Statistics Table (Page 2-8) provides ozone summary statistics for various indices computed on a monthly basis for an entire year. Growing season (generically defined to be May 1 - September 30) and annual statistics are also presented under the "MAY-SEP" and "ANNUAL" columns, respectively. All concentrations are expressed in the units of parts per billion (PPB) and exposures in parts per billion-hours (PPB-HR). The definitions for each of the statistics appearing on the Quick Look Annual Summary Table are given below.


- (1) **Daily 1-Hr Maximum.** The maximum 1-hour average concentration recorded during each month, the growing season or the year regardless of the number of valid hourly observations recorded during a given day. The number in parentheses below this statistic, (N), indicates the number of days in the month, growing season, or year with valid data.
- (2) **Average Daily Maximum.** The average of all Daily 1-Hr Maxima during the month regardless of the number of Daily 1-Hr Maxima recorded during the month. For the "MAY-SEP" column the average of all the Daily Maxima recorded during the growing season is given. For the "ANNUAL" column the average of all the Daily Maxima is given. N is as in (1) above.
- (3) **Maximum Daily Mean.** The maximum of the valid daily means computed for each month, the growing season ("MAY-SEP" column), and the year ("ANNUAL" column). A valid daily mean is one for which 75% of the observations are available for each day, i.e., 18 hours. N is the number of days during each month, growing season, and year with at least 18 observations.
- (4) **Average Daily Mean.** The average of all valid daily means for the month, the growing season ("MAY-SEP" column), and the year ("ANNUAL" column). N is as in (3) above.
- (5) **Max Peak:Min Ratio.** The ratio of the Daily 1-Hr Maximum to the Daily 1-Hr Minimum. A ratio is computed only if a valid Daily Mean is computed and if the Daily 1-Hr Minimum is not equal to zero. N is the number of days with a valid Peak:Min ratio.
- (6) **Average Peak:Min Ratio.** The average of all Peak:Min ratios for the month, growing season, or year. N is as in (5) above.
- (7) **Max 9AM-4PM Average.** The maximum of all valid 9AM-4PM Averages computed for the month, growing season, or year. A valid 9AM-4PM Average is one which has 75% of the observations available during that time period (i.e., 6 hours. N is the number of days with valid averages.)

- (8) **Monthly 9AM-4PM Average.** The average of all valid 9AM-4PM Averages for the month, growing season, or year. N is as in (7) above.
- (9) **Max 7AM-7PM Average.** The maximum of all valid 7AM-7PM Averages computed for the month, growing season, or year. A valid 7AM-7PM Average is one which has 75% of the observations available during that time period, i.e., 9 hours. N is the number of days with valid averages.
- (10) **Monthly 7AM-7PM Average.** The average of all valid 7AM-7PM averages for the month, growing season, or year. N is as in (9) above.
- (11) **Monthly Mean.** The average of all 1-Hr ozone concentrations recorded during the month, growing season, or year. A mean is computed regardless of the number of hours with valid data. N is the number of hours with valid observations.
- (12) **SUM0 Exposure Index.** The monthly sum of all hourly ozone concentrations. Units are PPB-HR. The "MAY-SEP" column sums across the months of May through September to give the cumulative exposure for the growing season. The "ANNUAL" column sums across every month to give the cumulative exposure for the year. N is the number of hours with valid observations and is the same N as in (11) above.
- (13) **SUM60 Exposure Index.** The monthly sum of all hourly ozone concentrations equaling or exceeding 60 PPB. Units are PPB-HR. The "MAY-SEP" column sums across the months of May through September to give the cumulative exposure for the growing season. The "ANNUAL" column sums across every month to give the cumulative exposure for the year. N is the number of hours equaling or exceeding 60 PPB during the month, growing season, or year.
- (14) **SUM80 Exposure Index.** The monthly sum of all hourly ozone concentrations equaling or exceeding 80 PPB. Units are PPB-HR. The "MAY-SEP" column sums across the months of May through September to give the cumulative exposure for the growing season. The "ANNUAL" column sums across every month to give the cumulative exposure for the year. N is the number of hours equaling or exceeding 80 PPB during the month, growing season, or year.
- (15) **W126 Exposure Index.** The monthly sum of all hourly ozone concentrations where each concentration is weighted by a function that gives greater emphasis to the higher hourly concentrations while still including the lower ones. This weighting function provides a weighting value that is unique for each hourly ozone concentration. The weighting function, as described by Lefohn, Laurence, and Kohut¹ is:

$$w_i = \frac{1}{1 + 4403 \exp(-.126c_i)}$$

where

Weighting Function Used To Calculate W126 Exposure Index



w_i = weighting value for hourly concentration i ,
and
 c_i = hourly concentration i in PPB.

The graph of weighting value versus ozone concentration, in the figure to the left, illustrates the greater weights given to higher hourly ozone concentrations.

Each hour's weighting value is multiplied by its corresponding hourly concentration. This product is summed over all the valid hours in each month to calculate the monthly W126 exposure.

Thus, the monthly W126 exposure is:

$$W126 = \sum_{i=1}^n w_i c_i$$

where

- W126 = monthly W126 exposure index,
- w_i = weighting value for hourly concentration i ,
- c_i = hourly concentration i in PPB, and
- n = number of hours in the month with valid ozone concentrations.

The "MAY-SEP" column sums across the months of May through September to give the cumulative exposure for the growing season. The "ANNUAL" column sums across every month to give the cumulative exposure for the year. The exposure units are PPB-HR.

Because each hour contributes to this exposure index, N is the number of hours with valid observations and is the same N as in (11) and (12) above.

The U.S. Environmental Protection Agency usually considers air quality statistics, such as a mean, to be "valid" (i.e., representative of the parameter being estimated for the time interval in question) only if 75% or more of the total possible observations have been measured during that time interval. Therefore, one should exercise caution when comparing these statistics between months and sites, particularly those that are not averages (e.g., maxima and exposures) whenever the number of valid observations is less than 75% of the total possible.

References

1. Lefohn, A.S., J. A. Laurence, and R. J. Kohut. 1988. A Comparison of Indices That Describe the Relationship Between Exposure to Ozone and Reduction in the Yield of Agricultural Crops. *Atmospheric Environment* 22, 1229-1240.

4.2 AIR QUALITY GLOSSARY

Acid Deposition: Air pollution produced when acid chemicals are incorporated into rain, snow, fog, or mist.

Aerometric Information Retrieval System (AIRS): A computer-based database of U.S. air pollution information administered by the EPA Office of Air Quality Planning and Standards (U.S. Environmental Protection Agency).

AIRWeb: Air Resources Web, an air quality information retrieval system for U.S. parks and wildlife refuges developed by the Air Resources Division of the National Park Service and the Air Quality Branch of the Fish and Wildlife Service.

Air Pollutant: An unwanted chemical or other material found in the air.

Air Pollution: Degradation of air quality resulting from unwanted chemicals or other materials occurring in the air.

Air Quality: The properties and degree of purity of air to which people and natural and heritage resources are exposed (in the context of national parks).

Air Pollution Control Permitting Process: Process by which facilities are permitted to emit specified types and quantities of air pollutants.

Air Quality Related Values (AQRVs): Values including visibility, flora, fauna, cultural and historical resources, odor, soil, water, and virtually all resources that are dependent upon and affected by air quality. "These values include visibility and those scenic, cultural, biological, and recreation resources of an area that are affected by air quality." (43 Fed. Reg. 15016)

Ambient Air: Air that is accessible to the public.

Class I: Areas of the country set aside under the Clean Air Act to receive the most stringent degree of air quality protection.

Class II: Areas of the country protected under the Clean Air Act but identified for somewhat less stringent protection from air pollution damage than Class I, except in specified cases.

Clean Air Act: Originally passed in 1963, our current national air pollution control program is based on the 1970 version of the law. Substantial revisions were made by the 1990 Clean Air Act Amendments.

Continuous Sampling Device: An air analyzer that measures air quality components continuously.

Criteria: Information on health and/or environmental effects of pollution (in the context of criteria air pollutants).

Criteria Air Pollutant: A group of very common air pollutants regulated by EPA on the basis of criteria and for which a National Ambient Air Quality Standard is established (SO₂, NO₂, PM₁₀, Pb, CO, O₃).

Emissions: Release of pollutants into the air from a source.

Environmental Protection Agency (EPA): The federal agency responsible for regulating air quality.

Monitoring: Measurement of air pollution.

National Ambient Air Quality Standards (NAAQS): Permissible levels of criteria air pollutant established to protect public health and welfare.

Ozone (O₃): A criteria air pollutant that is a strong oxidizing agent, reactive with many other compounds and surfaces, and a health hazard in high concentrations. Ozone is formed by nitrogen oxides and organic compounds reacting in sunlight.

Source: Any place or object from which air pollutants are released. Sources that are fixed in space are stationary sources; sources that move are mobile sources.

Sulfur Dioxide (SO₂): A criteria air pollutant that is a gas produced by burning coal and some industrial processes.

* Recent updates to this glossary may be found on the NPSARD AIRWeb - <http://www.aqd.nps.gov/natnet/ard/glossary.htm>.

4.3 GLOSSARY OF AIR QUALITY UNITS

| Units Conversion Table | | | |
|---|--|-----------|---|
| Parameter Type | Multiply | By | To Obtain |
| Pollutant | ppm | 1000 | ppb |
| | ppm | 1960 | $\mu\text{g}/\text{m}^3$ Ozone (at 25°C) |
| | ppm | 2615 | $\mu\text{g}/\text{m}^3$ Sulfur Dioxide (at 25°C) |
| | ppb | 0.001 | ppm |
| | ppb | 1.960 | $\mu\text{g}/\text{m}^3$ Ozone (at 25°C) |
| | ppb | 2.615 | $\mu\text{g}/\text{m}^3$ Sulfur Dioxide (at 25°C) |
| | $\mu\text{g}/\text{m}^3$ Ozone (25°C) | 0.0005102 | ppm |
| | $\mu\text{g}/\text{m}^3$ Ozone (25°C) | 0.5102 | ppb |
| | $\mu\text{g}/\text{m}^3$ Sulfur Dioxide (25°C) | 0.0003824 | ppm |
| | $\mu\text{g}/\text{m}^3$ Sulfur Dioxide (25°C) | 0.3824 | ppb |
| | | | |
| Wind Speed | m/s | 2.05 | mph |
| | mph | 0.489 | m/s |
| | | | |
| Solar Radiation | ly/min | 697 | w/m^2 |
| | w/m^2 | 0.00143 | ly/min |
| | | | |
| Precipitation | mm/hr | 0.0394 | in/hr |
| | in/hr | 25.4 | mm/hr |
| | | | |
| Temperature | $^{\circ}\text{C} + 17.78$ | 1.8 | $^{\circ}\text{F}$ |
| | $^{\circ}\text{F} - 32$ | 5/9 | $^{\circ}\text{C}$ |
| <p>Where:</p> <p>ppm = parts per million</p> <p>ppb = parts per billion</p> <p>$\mu\text{g}/\text{m}^3$ = micrograms per cubic meter (at 25°C)</p> <p>m/s = meters per second</p> <p>mps = miles per hour</p> <p>ly/min = langley's per minute</p> <p>w/m^2 = watts per square meter</p> <p>mm/hr = millimeters per hour</p> <p>in/hr = inches per hour</p> <p>$^{\circ}\text{C}$ = degrees centigrade</p> <p>$^{\circ}\text{F}$ = degrees fahrenheit</p> | | | |