


Annual Data Summary
ACADIA NATIONAL PARK
2002
National Park Service
Gaseous Air Pollutant Monitoring Network



AIR RESOURCES DIVISION
RESEARCH AND MONITORING BRANCH
12795 West Alameda Parkway
P.O. Box 25287
Lakewood, Colorado 80225
Telephone: (303) 969-2820
Fax: (303) 969-2822



The state of Maine and the National Park Service maintained two air pollutant monitoring sites in Acadia National Park during 2002. Data from each site are summarized in this report.

Annual Data Summary
ACADIA NATIONAL PARK
Cadillac Mountain
2002
National Park Service
Gaseous Air Pollutant Monitoring Network



AIR RESOURCES DIVISION
RESEARCH AND MONITORING BRANCH
12795 West Alameda Parkway
P.O. Box 25287
Lakewood, Colorado 80225
Telephone: (303) 969-2820
Fax: (303) 969-2822

This Annual Data Summary was prepared under NPS Contract C2350010840 by:

Air Resource Specialists, Inc.
1901 Sharp Point Drive, Suite E
Fort Collins, Colorado 80525
Telephone: (970) 484-7941
Fax: (970) 484-3423

For additional copies of this report or reports for other NPS units, contact:

National Park Service Air Resources Division
Information Management Center
c/o Air Resource Specialists, Inc.
1901 Sharp Point Drive, Suite E
Fort Collins, Colorado 80525
Telephone: (970) 484-7941
Fax: (970) 484-3423
E-Mail: AIR-IMC@AIR-RESOURCE.COM

or

National Park Service
Air Resources Division
P.O. Box 25287
Lakewood, Colorado 80225-02587
Telephone: (303) 969-2130
E-Mail: AQ_INFO@AQD.NPS.GOV

ACKNOWLEDGEMENTS

The National Park Service Air Resources Division (ARD) recognizes the level of effort required by individual park units, site operators, auditors, cooperating state and local agencies, and ARD contractors. ARD sincerely appreciates the contributions of all participants in assisting with the collection, validation, and reporting of these air quality and meteorological data.

At Acadia National Park, the ARD specifically recognizes Bob Breen and Bill Gawley of the State of Maine Department of Environmental Protection Bureau of Air Quality for performing the technical and administrative skills required to help produce the data presented within this report.

TABLE OF CONTENTS

<u>Section</u>		<u>Page</u>
1.0	INTRODUCTION	1-1
1.1	The National Park Service Gaseous Pollutant Monitoring Network	1-1
1.2	Acadia National Park - Cadillac Mountain	1-3
2.0	DATA SUMMARY	2-1
2.1	Overview	2-1
2.2	Ozone Data Summary	2-7
2.3	Meteorological Data Summary	2-23
3.0	NATIONAL PARK SERVICE AIR RESOURCES DIVISION DATA SOURCES	3-1
4.0	GLOSSARY	4-1
4.1	Definitions and Computational Procedures for National Park Service Quick Look Annual Summary Statistics Table	4-1
4.2	Air Quality Glossary	4-4
4.3	Glossary of Air Quality Units	4-6

LIST OF FIGURES AND TABLES

	<u>Page</u>
Map of National Park Service Gaseous Pollutant Monitoring Network Ozone and Sulfur Dioxide Monitoring Sites	1-2
Site Specifications	1-4
Data Collection Statistics	2-2
First Quarter Data Stackplot	2-3
Second Quarter Data Stackplot	2-4
Third Quarter Data Stackplot	2-5
Fourth Quarter Data Stackplot	2-6
Ozone Quick Look Annual Summary Statistics	2-8
Ozone Annual Frequency Distribution	2-9
Ozone Daily 1-Hour Maximum Concentrations and National Ambient Air Quality Standards Comparison	2-10

LIST OF FIGURES AND TABLES (Continued)

	<u>Page</u>
Attainment Status with EPA Proposed Primary Ozone Standards	2-11
Ten Highest 1-Hour Average Ozone Concentrations	2-12
Episodes with 1-Hour Ozone Concentrations ≥ 100 ppb and > 124 ppb	2-13
Episodes with 8-Hour Ozone Concentrations >84 ppb	2-14
Ozone Rank Listings of Second Highest 1-Hour Average Concentrations, Maximum 8-Hour Average Concentrations, and Annual Sum60 Exposure Index for All NPS Monitoring Sites	2-15
Plot of Maximum Ozone Concentration Comparison for Three Years	2-16
Map of National Park Service Gaseous Pollutant Monitoring Network, Second Highest Hourly Ozone Concentration	2-17
Quarterly Diurnal Ozone Plots	2-18
Annual Diurnal Ozone Plot	2-19
Quarterly Ozone Pollutant Roses	2-20
Annual Ozone Pollutant Roses	2-21
Ozone Precision Check Data Summary	2-22
Summary of Selected Meteorological Data	2-24
Quarterly Wind Roses	2-25
Annual Wind Roses	2-26

1.0 INTRODUCTION

1.1 THE NATIONAL PARK SERVICE GASEOUS POLLUTANT MONITORING PROGRAM (GPMP)

Gaseous air pollutants, including ozone and sulfur dioxide, are of concern to the National Park Service (NPS). Pollutants like these can affect park unit biological resources as well as the health of park unit residents and visitors. The NPS established a gaseous pollutant monitoring program for several pollutants linked to effects on NPS resources. This program was designed to meet certain resource management objectives.

The primary objective of this monitoring program is to establish the status and trends of park unit air quality conditions and to determine if a park unit is exceeding the National Ambient Air Quality Standards established by the U.S. Environmental Protection Agency (EPA) to protect public health and welfare. In addition, such monitoring is designed to detect changes or trends in pollution levels over time. A monitoring station may also be established if there is documented biological injury due to air pollution in a park unit. Information on ambient air pollution levels is an important part of research on effects of air pollutants on NPS resources, and can help confirm suspected causes of observed effects.


Other monitoring objectives call for the collection of data to support the National Park Service's required involvement in both the development of state air quality control plans, and the evaluation of permit applications for new or expanding air pollution sources wishing to locate near park units. The Clean Air Act gives federal land managers and superintendents an affirmative responsibility to protect air quality related values in Class I areas and to assess whether new sources will have an adverse impact on park unit resources and values. Information on air quality levels in NPS units can also be used to evaluate the performance of atmospheric models that simulate how pollutants are transported into park units and predict impacts on the park unit caused by air pollution sources.

The National Park Service Gaseous Pollutant Monitoring Program site locations and measured parameters collected in this reporting year are shown on the map on the following page. During this reporting period, 47 monitoring sites in 37 units of the National Park System had some combination of ozone, sulfur dioxide, nitrogen, meteorological, and Clean Air Status and Trends Network (CASTNet) dry deposition monitoring. Monitoring methods and quality assurance procedures used in the national park network meet the applicable 40 CFR Part 58 EPA requirements. This allows for the direct comparison of NPS collected data with that collected by the EPA, and state and local air pollution control agencies. Data collected by this network are incorporated in the EPA Aerometric Information Retrieval System (AIRS) database which is a national database of all air quality data collected throughout the country. These data are also stored in the NPS Air Resources Division's Information Management Center (IMC) that allows for easy access and analysis of data.

This report includes a variety of data summaries for data collected at an individual monitoring site at a national park unit during this reporting period. These summaries highlight the average range and frequency of the data collected during the year. A digital copy of all data collected during the year and data summary products are available; see Section 3.0 for information on obtaining these data. Individual reports are generated for each site where monitoring was conducted in the national park network.

NATIONAL PARK SERVICE GASEOUS POLLUTANT MONITORING NETWORK

2002 Monitoring Sites



1.2 ACADIA NATIONAL PARK

Acadia National Park, a Class I area, is located along the coast of Maine in Knox and Hancock Counties. Its location and site specifications are presented on the next page.

The mandate from the United States Congress states Acadia National Park was established "...to conserve the scenery and the natural and historic objects and the wildlife therein, and to provide for the enjoyment of the same in such a manner and by such means as will leave them unimpaired for the enjoyment of the future generations." The purpose of Acadia is to perpetuate the historical traditions and natural history of a portion of the Maine coastline and to offer visitors an increased awareness of their American heritage.

Sieur de Monts National Monument was established by Presidential Proclamation in 1916. In 1919, the name was changed to Lafayette National Park and again in 1929 to Acadia National Park.

Acadia National Park conserves panoramas and scenic vistas of mountains, rocky coasts, islands and the ocean in a combination that is unequalled along the Atlantic coast. When viewed from seaside cliffs, islands along the coast spread out to the horizon. Behind the rocky shore, the rounded summits of glaciated mountains rise to 1,530 feet. Throughout the mountains are small clear lakes surrounded by spruce, fir and hardwoods. A major concern is residential and commercial development on private lands which threatens to encroach on park vistas offering views of forests, cliffs, and oceans.

The forests of Mount Desert Island have a special quality. The northern coniferous and temperate deciduous forests meet and overlap, bringing to the island an exceptional variety of trees, shrubs and herbaceous plant life. About 1,500 species of plants are found in Acadia, but the majority of these cannot be considered common. For example, scrub oak is only found on Acadia Mountain and a few nearby mountain slopes.


Recent research has found a number of native plants including broad-leaf aster (*Aster macrophyllus*), spreading dogbane (*Androsaemifolium*), black cherry (*Prunus serotina*), and white ash (*Fraxinus americana*) to be sensitive to ambient levels of ozone. Field surveys have confirmed the presence of foliar injury to some of these species growing under natural conditions at Acadia. Other research has indicated a positive correlation between ozone concentration and reduced tree-ring growth in several white pine stands in the park. There have also been a number of other biotic and abiotic agents identified that impact park vegetation including spruce budworm, gypsy moths, insects, and a fungus that produces needle injury to white pine.

Approximately 50 species of mammals and 327 species of birds have been reported in Acadia National Park, as well as amphibians and reptiles. The dominant mammals include raccoon, mink, skunk, red fox, harbor seal, beaver, snowshoe hare, white-tailed deer and harbor porpoise.


In addition to the issue of surrounding growth and development which encroaches primarily on water and visual resources, there are a number of air pollution related concerns. Acid deposition, ozone, mercury and other toxins including PCB's have all been identified as impacts to Acadia's flora, fauna, and hydrogeochemical resources.

Acadia National Park Cadillac Mountain

Ozone Monitoring
NPS Gaseous Air Pollutant Monitoring Network



Air Quality Station



	SITE IDENTIFICATION	MAP INFORMATION
	Site Abbreviation: ACAD-CM AIRS ID NO.: 29-009-0102	Mean Elevation: 466 m Longitude: 68°13'40"W Latitude: 44°20'50"N UTM Zone: 19
	INSTRUMENTATION	
	O ₃ Analyzer Wind Direction Relative Humidity Temperature Wind Speed Calibrator	Easting: 561551m Northing: 4910512m Map Reference: Bar Harbor 44068-A1 1:100,000

2.0 DATA SUMMARY

2.1 OVERVIEW

Based on the site specifications during this annual reporting period, data summaries and statistics are provided in this section.

Data Collection Statistics
Acadia National Park
Cadillac Mountain

Final Validation

01/01/2002 - 12/31/2002

Parameter	Interval	Par Code	Data Recovery			Valid Data	
			No. Possible	No. Collected	% Collected	No. Valid	% Valid
Ozone Analyzer	hourly	O3	6566	6362	96.9	6353	96.8
Scalar Wind Speed	hourly	SWS	3372	3371	100.0	3057	90.7
Vector Wind Speed	hourly	VWS	3336	3335	100.0	3011	90.3
Vector Wind Direction	hourly	VWD	3336	3335	100.0	3011	90.3
Scalar Wind Direction	hourly	SWD	3336	3335	100.0	3011	90.3
Standard Deviation for Wind Direction	hourly	SDWD	3336	3335	100.0	3010	90.2
Ambient Temperature (aspirated)	hourly	TMP	3374	3373	100.0	3343	99.1
Relative Humidity	hourly	RH	3374	3373	100.0	3370	99.9

Notes: The percent valid is calculated against the number possible. Automatic zeros and spans are performed daily on most ambient gas analyzers, therefore, no ambient data can be collected during this time. As a result, the maximum percent valid for ambient gas data typically can not be greater than 95.8.

Performance Goals:


Quarterly Criteria:

100% of sites, >= 85% valid data capture
90% of sites, >= 90% valid data capture
80% of sites, >= 95% valid data capture

Monthly Criteria:

100% of sites, >= 60% valid data capture
90% of sites, >= 75% valid data capture
80% of sites, >= 85% valid data capture


Acadia National Park - Cadillac Mountain



Final Validation

First Quarter 2002


Acadia National Park - Cadillac Mountain



Final Validation

Second Quarter 2002


Acadia National Park - Cadillac Mountain



Final Validation

Third Quarter 2002

Acadia National Park - Cadillac Mountain



Final Validation

Fourth Quarter 2002

2.2 OZONE DATA SUMMARY

Ozone Quick Look Annual Summary Statistics
Acadia National Park
Cadillac Mountain
01/01/2002 - 12/31/2002

STATISTIC	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC	MAY- SEP	ANNUAL
DAILY 1-HR MAXIMUM	70	141	99	127	127	127	127	127	127	68	48	50	141	141
NO. OF DAYS	(29)	(31)	(30)	(31)	(31)	(30)	(31)	(31)	(30)	(31)	(30)	(31)	(153)	(274)
AVERAGE DAILY MAXIMUM	50	57	56	63	51	39	37	39	39	39	37	39	57	50
NO. OF DAYS	(29)	(31)	(30)	(31)	(30)	(31)	(30)	(31)	(30)	(31)	(30)	(31)	(153)	(274)
MAXIMUM DAILY MEAN	52	70	68	86	77	53	40	45	45	45	40	45	86	86
NO. OF DAYS	(28)	(31)	(30)	(31)	(30)	(31)	(30)	(31)	(30)	(31)	(30)	(31)	(153)	(273)
AVERAGE DAILY MEAN	44	46	43	45	38	33	31	32	32	33	31	32	43	39
NO. OF DAYS	(28)	(31)	(30)	(31)	(30)	(31)	(30)	(31)	(30)	(31)	(30)	(31)	(153)	(273)
MAX PEAK:MIN RATIO	1.892	5.143	2.820	6.188	3.300	4.529	2.500	4.875	45.000	45.000	4.875	45.000	6.188	45.000
NO. OF DAYS	(28)	(31)	(30)	(31)	(30)	(31)	(30)	(31)	(30)	(31)	(30)	(31)	(153)	(273)
AVERAGE PEAK:MIN RATIO	1.333	1.630	1.649	2.051	1.841	1.551	1.893	2.988	2.988	2.988	1.893	2.988	1.827	1.884
NO. OF DAYS	(28)	(31)	(30)	(31)	(30)	(31)	(30)	(31)	(30)	(31)	(30)	(31)	(153)	(273)
MAX 9AM-4PM AVERAGE	56	79	83	81	73	61	45	45	45	61	45	45	83	83
NO. OF DAYS	(26)	(31)	(30)	(31)	(30)	(31)	(30)	(31)	(30)	(31)	(30)	(31)	(152)	(270)
MONTHLY 9AM-4PM AVERAGE	45	47	43	42	39	34	31	34	34	34	31	34	43	40
NO. OF DAYS	(26)	(31)	(30)	(31)	(30)	(31)	(30)	(31)	(30)	(31)	(30)	(31)	(152)	(270)
MAX 7AM-7PM AVERAGE	54	78	85	72	84	61	43	45	45	61	43	45	85	85
NO. OF DAYS	(28)	(31)	(30)	(31)	(30)	(31)	(30)	(31)	(30)	(31)	(30)	(31)	(153)	(273)
MONTHLY 7AM-7PM AVERAGE	45	47	43	41	44	39	34	33	33	34	31	33	43	40
NO. OF DAYS	(28)	(31)	(30)	(31)	(30)	(31)	(30)	(31)	(30)	(31)	(30)	(31)	(153)	(273)
MONTHLY MEAN	44	46	43	40	45	38	33	33	33	33	31	33	43	39
NO. OF HOURS	(646)	(710)	(696)	(721)	(723)	(695)	(724)	(705)	(733)	(733)	(705)	(733)	(3545)	(6353)
SUM0 EXPOSURE INDEX	28446	32999	30030	29056	32374	26577	23719	21722	23865	23865	21722	23865	151036	248788
NO. OF HOURS	(646)	(710)	(696)	(721)	(723)	(695)	(724)	(705)	(733)	(733)	(705)	(733)	(3545)	(6353)
SUM60 EXPOSURE INDEX	597	6971	5686	5897	14094	5429	686	-	38077	39360	-	-	38077	39360
NO. OF HOURS	(9)	(105)	(75)	(80)	(175)	(74)	(11)	(0)	(509)	(529)	(0)	(0)	(509)	(529)
SUM80 EXPOSURE INDEX	-	431	1887	2035	7466	1220	-	-	13039	13039	-	-	13039	13039
NO. OF HOURS	(0)	(5)	(20)	(23)	(80)	(13)	(0)	(0)	(141)	(141)	(0)	(0)	(141)	(141)
W126 EXPOSURE INDEX	2198	5639	5287	5296	11898	4514	884	386	36627	36627	386	525	32633	36627
NO. OF HOURS	(646)	(710)	(696)	(721)	(723)	(695)	(724)	(705)	(733)	(733)	(705)	(733)	(3545)	(6353)

Concentrations in parts per billion (ppb)
Exposures in parts per billion-hours (ppb-hr)
Final Validation

* Statistics defined in the Quick Look subsection of the Glossary

Frequency Distribution															
Acadia National Park Cadillac Mountain															
Monitoring Season: 04/01/02 - 09/30/02 ¹															
Averaging Period	% Obs. ³	# Obs. ²	Min. Obs. ⁴	10	30	50	Percentile ⁵			99	Max. Obs.	2nd Max.	Arith. Mean	Geo. Mean	Geo. Stdv.
							70	90	95						
1-Hour	99	4191	0.024	0.036	0.044	0.050	0.058	0.088	0.099	0.127	0.141	0.127	0.0558	0.0524	1.41
Concentrations in parts per million (ppm)															

¹Records for this report are selected in accordance with the AIRS Geo-Common file criteria. These criteria are based on the state-specific Monitoring Season defined in AIRS.

²The number of observations (# Obs.) includes all valid observations recorded within the Monitoring Season.

³The percent of valid observations (% Obs.) is the percentage of valid days to the number of possible monitoring days during the Monitoring Season. A valid day is defined as a day with 9 or more valid observations between 9:00 a.m. and 9:00 p.m..

⁴The minimum observation value (Min. Obs.) is the minimum daily maximum recorded during the Monitoring Season.

⁵The percentiles and other statistics are derived from the daily maximums.

Ozone Standards Report and
Daily Maximum 1-Hour Concentrations (ppm)

Acadia National Park
Cadillac Mountain

01/01/2002 - 12/31/2002

Day	Jan-02	Feb-02	Mar-02	Apr-02	May-02	Jun-02	Jul-02	Aug-02	Sep-02	Oct-02	Nov-02	Dec-02									
1	T	F	F	M	.047	W	.064	S	.080	M	.035	T	.036	S	.068	T	.037	F	.033	S	
2	W	S	S	T	.052	T	.051	S	.092	T	.029	F	.038	M	.061	W	.037	S	.040	M	
3	T	S	S	.059	W	.052	F	.039	M	.058	W	.048	S	.048	T	.045	T	.037	S	.038	T
4	F	M	M	.044	T	.056	S	.046	T	.095	T	.099	S	.043	W	.040	F	.036	M	.040	W
5	S	T	T	.046	F	.072	S	.055	W	.053	F	.088	M	.040	T	.048	S	.038	T	.039	T
6	S	W	W	.044	S	.077	M	.047	T	.031	S	.038	T	.052	F	.036	S	.040	W	.037	F
7	M	T	T	.049	S	.062	T	.041	F	.032	S	.031	W	.054	S	.033	M	.033	T	.035	S
8	T	F	F	.053	M	.050	W	.043	S	.082	M	.037	T	.088	S	.037	T	.038	F	.038	S
9	W	S	S	.054	T	.049	T	.058	S	.051	T	.046	F	.083	M	.039	W	.045	S	.039	M
10	T	S	S	.048	W	.065	F	.042	M	.042	W	.075	S	.127*	T	.038	T	.048	S	.039	T
11	F	M	M	.052	T	.053	S	.046	T	.032	T	.107	S	.077	W	.040	F	.043	M	.031	W
12	S	T	T	.053	S	.039	W	.058	F	.058	F	.117	M	.036	T	.041	S	.043	T	.040	T
13	S	W	W	.049	S	.053	M	.044	T	.093	S	.118	T	.046	F	.040	S	.034	W	.050	F
14	M	T	T	.043	S	.053	T	.056	F	.097	S	.127*	W	.050	S	.039	M	.032	T	.045	S
15	T	F	F	.047	M	.044	W	.047	S	.061	M	.119	T	.051	S	.032	T	.031	F	.034	S
16	W	S	S	.070	T	.070	T	.033	S	.054	T	.087	F	.041	M	.045	W	.034	S	.039	M
17	T	S	S	.056	W	.063	F	.042	M	.052	W	.065	S	.044	T	.042	T	.038	S	.039	T
18	F	M	M	.046	T	.036	S	.051	T	.052	T	.094	S	.038	W	.031	F	.037	M	.041	W
19	S	T	T	.064	F	.036	S	.039	W	.049	F	.072	M	.081	T	.042	S	.042	T	.049	T
20	S	W	W	.044	S	.046	M	.049	T	.052	S	.037	T	.060	F	.029	S	.039	W	.045	F
21	M	T	T	.052	S	.041	T	.101	F	.032	S	.037	W	.061	S	.033	M	.039	T	.045	S
22	T	F	F	.042	M	.055	W	.081	S	.062	M	.065	T	.044	S	.034	T	.037	F	.029	S
23	W	S	S	.044	T	.070	T	.078	S	.099	T	.058	F	.036	M	.029	W	.036	S	.031	M
24	T	S	S	.047	W	.092	F	.052	M	.029	W	.037	S	.024	T	.026	T	.032	S	.034	T
25	F	M	M	.052	T	.044	S	.079	T	.036	T	.045	S	.038	W	.037	F	.030	M	.032	W
26	S	T	T	.048	F	.046	S	.141*	W	.035	F	.074	M	.032	T	.046	S	.033	T	.038	T
27	S	W	W	.050	S	.062	M	.094	T	.032	S	.046	T	.049	F	.042	S	.037	W	.043	F
28	M	T	T	.053	S	.064	T	.056	F	.049	S	.024	W	.040	S	.031	M	.040	T	.044	S
29	T		F	.050	M	.070	W	.038	S	.050	M	.026	T	.027	S	.037	T	.040	F	.034	S
30	W		S	.050	T	.055	T	.054	S	.058	T	.036	F	.044	M	.037	W	.036	S	.038	M
31	T		S		.072	F	.046	W	.035	S					.036	T					
Valid Days	0	0	0	27	31	30	31	31	30	30	31	30	30	30	31	30	30	30	30	30	30
Maximum	.000	.000	.000	.070	.092	.141	.099	.127	.127	.068	.048	.050									
Violations	0	0	0	0	0	1	0	1	1	0	0	0	0	1	0	0	0	0	0	0	0

6329 Total Samples 3 Daily-maxima exceeding the standard of .12 ppm (starred[*])
72.4 % Possible 1 Missing days assumed to be less than the standard
271 Valid daily maxima 0 Daily_maximas exceed the alert level of .200 ppm
Concentrations in parts per million (ppm)
Final Validation 4/22/03

Acadia National Park
 Cadillac Mountain
 2002 Attainment Status With U.S. Environmental Protection Agency (EPA)
 PRIMARY Ozone National Ambient Air Quality Standard

Ozone Season: April through September

The primary National Ambient Air Quality Standard for ozone is designed to protect human health. The level of the primary ozone standard promulgated by the EPA on July 18, 1997 is 0.08 parts per million (ppm) [80 parts per billion, (ppb)], daily maximum 8-hour average. The primary ozone standard is met at an ambient monitoring site when the 3-year average of the annual fourth-highest daily maximum 8-hour average ozone concentration is less than or equal to 0.08 ppm. This standard is not met when the 3-year average is greater than 0.08 ppm. Using the EPA's rounding convention, a computed 3-year average ozone concentration of 0.085 ppm (85 ppb) is the smallest value that is greater than the level of the 0.08 ppm standard.

The primary standard requires 90 percent data completeness, on average, during the 3-year period, with no single year within the period having less than 75 percent data completeness. This data completeness requirement would have to be satisfied in order to determine that the standard has been met at a monitoring site. However, calendar years with less than 75 percent data completeness are included in the computation if the annual fourth-highest daily maximum 8-hour concentration is greater than the level of the standard. A site could be found not to have met the standard with less than complete data. The percent data completeness is the percent of valid ozone monitoring days. A day is valid if valid 8-hour averages are available for at least 75 percent of possible hours in the day (i.e., at least 18 of the 24 averages). An 8-hour average is considered valid if at least 75 percent (or 6) of the hourly averages for the 8-hour period are available.

The table below lists the 3-year average fourth-highest daily maximum 8-hour ozone concentration based on data collected during the reported year and the two previous years. This is the number to compare to the level of the new primary standard. The 3-year average data completeness percent and the reported year highest five daily maximum 8-hour averages are also tabulated. A 'No' in the Data Comp % Met? column indicates EPA data completeness requirement was not met for the three-year period.

Year	3-Year Avg 4th High Daily Max 8-hr Ozone (ppb)	3-Year Avg Data Complete %	Data Complete % Met?	Annual 1st High Daily Max 8-hr Ozone (ppb)	Annual 2nd High Daily Max 8-hr Ozone (ppb)	Annual 3rd High Daily Max 8-hr Ozone (ppb)	Annual 4th High Daily Max 8-hr Ozone (ppb)	Annual 5th High Daily Max 8-hr Ozone (ppb)
2002	92	93%	Yes	111	106	106	100	99

Ozone Analyzer			
10 Highest Daily 1-Hour Average Maximum Concentrations			
Acadia National Park			
Cadillac Mountain			
Final Validation			
01/01/2002 - 12/31/2002			
Value	Date	Hour	Concentration (ppm)
Ozone Analyzer			
1	06/26/2002	19	0.141
2	08/14/2002	22	0.127
3	09/10/2002	17	0.127
4	08/15/2002	2	0.119
5	08/13/2002	21	0.118
6	08/12/2002	22	0.117
7	08/11/2002	15	0.107
8	06/21/2002	18	0.101
9	07/23/2002	14	0.099
10	08/04/2002	20	0.099**

** This value was also recorded on one or more days later in the reported period.

Episodes with 1-Hour Ozone Concentrations
 ≥ 100 ppb and > 124 ppb

Acadia National Park
 Cadillac Mountain

01/01/2002 - 12/31/2002

FINAL VALIDATION

Site	Date	Beginning Hour	No. Hours		Max (ppb)
			≥ 100 ppb	>124 ppb	
ACAD-CM	06/21/02	18	1	0	101
ACAD-CM	06/26/02	18	2	1	141
ACAD-CM	08/11/02	15	1	0	107
ACAD-CM	08/12/02	18	5	0	117
ACAD-CM	08/12/02	23	1	0	110
ACAD-CM	08/13/02	0	4	0	106
ACAD-CM	08/13/02	19	4	0	118
ACAD-CM	08/13/02	23	1	0	116
ACAD-CM	08/14/02	0	2	0	109
ACAD-CM	08/14/02	21	2	1	127
ACAD-CM	08/14/02	23	1	0	112
ACAD-CM	08/15/02	1	4	0	119
ACAD-CM	09/10/02	17	3	1	127
		Total	31	3	141

Note: The primary and secondary national ambient air standard for ozone that applied in 1996 is 0.12 ppm over a one hour period not to be exceeded more than once per year. (A value greater than .12 ppm, 124 ppb, or 235 ug/m³ exceeds the standard.) (40 CFR 50.9 with reference to Appendix D and H.)

Episodes with 8-Hour Average Ozone Concentrations > 84 ppb

Acadia National Park

Cadillac Mountain

01/01/2002 - 12/31/2002

FINAL VALIDATION

Site	Date	Start and End Time of Daily Maximum 8-Hour Average > 84 ppb (hr)	Daily Maximum 8-Hour Average (ppb)	Number of 8-Hour Averages > 84 ppb During the Day
ACAD-CM	06/21/02	15 - 22	96	10
ACAD-CM	06/26/02	18 - 01	98	7
ACAD-CM	08/11/02	12 - 19	89	5
ACAD-CM	08/12/02	19 - 02	106	13
ACAD-CM	08/13/02	19 - 02	106	12
ACAD-CM	08/14/02	21 - 04	111	11
ACAD-CM	08/15/02	00 - 07	100	8
ACAD-CM	09/10/02	16 - 23	99	8
	8	Days with 8-hour average concentrations > 84 ppb		

Note: This table presents episodes of high ozone based on running 8-hour averages. In 1997, the EPA published new primary and secondary national ambient air quality standards for ozone based on 8-hour average ozone concentrations. Attainment of the new primary standard is reached if the annual fourth highest daily maximum 8-hour ozone concentration, averaged over three years, does not exceed 0.08 ppm (84 ppb or 157 ug/m³).



Ozone Rank Listings of Second Highest 1-Hour Average Concentrations, 4th Highest 8-Hour Average Concentrations, and Annual SUM60 Exposure Index for All NPS Monitoring Sites

01/01/2002 - 12/31/2002


Second Highest 1-Hour Average Concentration		
Site	Rank	Concentration (ppb)
ACAD-CM	1	127
CHAM-XX	2	127
JOTR-YV	3	127
SEKI-LK	4	126
SEKI-AS	5	124
SEKI-LP	6	124
GRSM-LR	7	122
CACO-XX	8	118
COWP-XX	9	118
ACAD-MH	10	117
GRSM-CM	11	117
GRSM-CD	12	115
COSW-BL	13	111
MACA-HM	14	110
PINN-ES	15	110
GRSM-CC	16	108
ROMO-LP	17	106
GRSM-PK	18	105
YOSE-TD	19	105
SHEN-BM	20	103
DEVA-PV	21	97
SAGU-PC	22	90
GRBA-MY	23	89
GRCA-AS	24	85
LAVO-ML	25	84
CHIR-ES	26	80
YOSE-MR	27	80
MEVE-MY	28	79
CHIS-XX	29	78
CANY-IS	30	77
CRMO-VC	31	75
YELL-WT	32	73
THRO-VC	33	71
MORA-TW	34	70
NOCA-MM	35	70
VOYA-SB	36	70
PEFO-HB	37	69
BIBE-KB	38	68
EVER-BC	39	68
DENA-HQ	40	65
GLAC-WG	41	59
VIIS-LP	42	57
HAVO-TH	43	50
OLYM-VC	44	44

4th Highest 8-hour Average Concentration		
Site	Rank	Concentration (ppb)
SEKI-LP	1	109
SEKI-LK	2	108
JOTR-YV	3	107
SEKI-AS	4	107
GRSM-CM	5	103
GRSM-LR	6	102
GRSM-CD	7	101
ACAD-CM	8	100
GRSM-PK	9	94
CACO-XX	10	93
COWP-XX	11	93
YOSE-TD	12	93
ACAD-MH	13	89
CHAM-XX	14	89
ROMO-LP	15	87
PINN-ES	16	86
SHEN-BM	17	86
MACA-HM	18	85
DEVA-PV	19	83
COSW-BL	20	82
GRSM-CC	21	82
GRCA-AS	22	79
SAGU-PC	23	77
LAVO-ML	24	75
GRBA-MY	25	74
CANY-IS	26	72
YOSE-MR	27	72
MEVE-MY	28	70
CHIR-ES	29	69
CRMO-VC	30	69
CHIS-XX	31	66
YELL-WT	32	66
VOYA-SB	33	65
BIBE-KB	34	62
THRO-VC	35	62
EVER-BC	36	57
DENA-HQ	37	55
PEFO-HB	38	55
GLAC-WG	39	52
MORA-TW	40	52
VIIS-LP	41	48
NOCA-MM	42	46
HAVO-TH	43	42
OLYM-VC	44	39


Annual Sum60 Exposure Index			
Site	Rank	Sum60 Count	
SEKI-LK	1	204306	2639
SEKI-AS	2	196849	2529
SEKI-LP	3	193795	2491
JOTR-YV	4	175177	2398
GRSM-CM	5	169849	2320
YOSE-TD	6	164764	2298
GRSM-LR	7	131936	1794
GRSM-CD	8	130649	1793
GRSM-PK	9	118538	1656
DEVA-PV	10	106174	1586
GRCA-AS	11	104360	1584
ROMO-LP	12	95145	1403
SHEN-BM	13	88006	1273
CANY-IS	14	68738	1068
COWP-XX	15	66858	924
PINN-ES	16	54322	767
MEVE-MY	17	49400	771
GRSM-CC	18	42779	611
CACO-XX	19	39974	546
MACA-HM	20	39775	571
ACAD-CM	21	39360	529
SAGU-PC	22	35867	540
COSW-BL	23	34655	491
CHAM-XX	24	34216	482
CRMO-VC	25	34186	537
CHIR-ES	26	32418	503
GRBA-MY	27	30461	466
YELL-WT	28	29522	470
ACAD-MH	29	26476	362
LAVO-ML	30	26432	394
YOSE-MR	31	22272	334
VOYA-SB	32	7405	116
BIBE-KB	33	7215	116
CHIS-XX	34	6974	106
THRO-VC	35	4004	63
DENA-HQ	36	996	16
PEFO-HB	37	951	15
EVER-BC	38	833	13
MORA-TW	39	453	7
NOCA-MM	40	267	4
GLAC-WG	41	124	2
HAVO-TH	42	0	0
OLYM-VC	43	0	0
VIIS-LP	44	0	0




NAAQS




Legend




FIRST QUARTER (JAN-MAR)




SECOND QUARTER (APR-JUN)




THIRD QUARTER (JUL-SEP)



FOURTH QUARTER (OCT-DEC)



—	NAAQS (0 ppm)	— —	Average of Valid O3 in ppm (Bar is equal to +/- one standard deviation)
—■—	Maximum of Valid O3 in ppm (□ Indicates Data Exceeds Plot Scale)	—▲—	Minimum of Valid O3 in ppm




Acadia National Park
Cadillac Mountain

Quarterly Ozone
Pollutant Rose


2002

FIRST QUARTER (JAN-MAR)




0.0% Collected 0.0% Valid
0 Possible /0 Collected /0 Valid
(includes WS and WD)

SECOND QUARTER (APR-JUN)




96.3% Collected 68.6% Valid
1128 Possible /1086 Collected /774 Valid
(includes WS and WD)

Ozone (ppb)




THIRD QUARTER (JUL-SEP)




96.9% Collected 96.9% Valid
2208 Possible /2139 Collected /2139 Valid
(includes WS and WD)

Final Validation

FOURTH QUARTER (OCT-DEC)



0.0% Collected 0.0% Valid
0 Possible /0 Collected /0 Valid
(includes WS and WD)



96.7% Collected 87.3% Valid
3336 Possible /3225 Collected /2913 Valid
(includes WS and WD)

Ozone Analyzer Precision Check Summary
Acadia National Park
Cadillac Mountain

Precision checks are required by the Environmental Protection Agency (EPA) of all monitoring instruments collecting data which are to be submitted to the EPA Aerometric Information Retrieval System (AIRS). A precision check is performed by challenging the pollutant analyzer with a known concentration of gas from the pollutant transfer standard. This precision check must be performed at least every 14 days of monitoring operation. The percent difference between the analyzer and the transfer standard is then calculated.¹ According to NPS Standard Operating Procedures, the pollutant analyzer must respond within 10% of the transfer standard. The table below gives the number of precision checks performed during each quarter, the average² of all the individual precision check percent differences for the quarter, and the upper and lower 95% probability limits³ for precision checks. The probability limits represent the interval having a 95% chance of containing the true average percent difference. The quarterly average percent difference and probability limits should ideally be within +/- 10%.

Final Validation 01/01/2002 - 12/31/2002				
Calendar Quarter	Number of Precision Checks	Average Percent Difference ^{1 2}	Lower 95% Probability Limit ³	Upper 95% Probability Limit ³
1	0			
2	11	0.42	-2.41	3.24
3	9	-0.61	-1.76	0.53
4	9	0.62	-0.53	1.76

¹ Percent Difference = $\frac{\text{analyzer} - \text{transfer std}}{\text{transfer std}} \times 100$.

² Average Percent Difference is the mean of all individual precision check percent differences during the quarter.

³ Upper/Lower 95% Probability Limits = (Average Percent Difference) +/- (1.96)(Standard Deviation of precision check percent differences in the quarter.)

2.3 METEOROLOGICAL DATA SUMMARY

Summary of Selected Meteorological Data

Acadia National Park

Cadillac Mountain

Final Validation

01/01/2002 - 12/31/2002


Parameter	Value	Units	Number	Std Dev
SCALAR WIND SPEED				
Average	6.2	m/s	3057	3.0
Maximum	23.0	m/s		
Percent calm = 0.00				
AMBIENT TEMPERATURE				
Average	15.3	degC	3343	5.5
Maximum	30.7	degC		
Minimum	0.4	degC		
RELATIVE HUMIDITY				
Average	79	percent	3370	21
Maximum	100	percent		
Minimum	28	percent		
	NA			
PRECIPITATION (Rainfall or Snow melt)				
Average non-zero rate	NA			
Maximum non-zero rate				
Minimum non-zero rate				
Accumulated during period				
SOLAR RADIATION				
Average Daily Total	NA			
Maximum Daily Total	NA			
Minimum Daily Total	NA			

Note: Calms are included in the average scalar wind speed and are defined as winds less than 0.5 m/s (1.0 mph).

Solar radiation terms are based on the calculation of the total amount of solar energy incident on a unit area during each day. The maximum and minimum daily totals are selected from the list of daily totals. The totals for all days are then added and divided by the number of days to yield the average daily total. Only days with 24 valid values are included in these statistics.


NA indicates instrument not available.

FIRST QUARTER (JAN-MAR)




0.0% Collected 0.0% Valid
0 Possible /0 Collected /0 Valid
(includes WS and WD)

SECOND QUARTER (APR-JUN)




99.9% Collected 71.3% Valid
1128 Possible /1127 Collected /804 Valid
(includes WS and WD)

Scalar Wind Speed (m/s)




THIRD QUARTER (JUL-SEP)




100.0% Collected 100.0% Valid
2208 Possible /2208 Collected /2207 Valid
(includes WS and WD)

Final Validation

FOURTH QUARTER (OCT-DEC)



0.0% Collected 0.0% Valid
0 Possible /0 Collected /0 Valid
(includes WS and WD)



3.0 NATIONAL PARK SERVICE AIR RESOURCES DIVISION DATA SOURCES

Meteorological and hourly gaseous data contained in this report may be obtained from the following sources:

- National Park Service AIRWeb (<http://www.aqd.nps.gov/natnet/ard>)
- EPA AIRS database
- Data requests directed to:

NPS Air Resources Division
Information Management Center
c/o Air Resource Specialists, Inc.
1901 Sharp Point Drive, Suite E
Fort Collins, Colorado 80525
Telephone: (970) 484-7941
Fax: (970) 484-3423
E-Mail: AIR-IMC@AIR-RESOURCE.COM

CASTNet concentration data may be obtained from the following Web site:

<http://www.epa.gov/castnet/data.html>

4.0 GLOSSARY

4.1 DEFINITIONS AND COMPUTATIONAL PROCEDURES FOR NATIONAL PARK SERVICE QUICK LOOK ANNUAL SUMMARY STATISTICS REPORT

The National Park Service Quick Look Annual Summary Statistics Table (Page 2-8) provides ozone summary statistics for various indices computed on a monthly basis for an entire year. Growing season (generically defined to be May 1 - September 30) and annual statistics are also presented under the "MAY-SEP" and "ANNUAL" columns, respectively. All concentrations are expressed in the units of parts per billion (PPB) and exposures in parts per billion-hours (PPB-HR). The definitions for each of the statistics appearing on the Quick Look Annual Summary Table are given below.


- (1) **Daily 1-Hr Maximum.** The maximum 1-hour average concentration recorded during each month, the growing season or the year regardless of the number of valid hourly observations recorded during a given day. The number in parentheses below this statistic, (N), indicates the number of days in the month, growing season, or year with valid data.
- (2) **Average Daily Maximum.** The average of all Daily 1-Hr Maxima during the month regardless of the number of Daily 1-Hr Maxima recorded during the month. For the "MAY-SEP" column the average of all the Daily Maxima recorded during the growing season is given. For the "ANNUAL" column the average of all the Daily Maxima is given. N is as in (1) above.
- (3) **Maximum Daily Mean.** The maximum of the valid daily means computed for each month, the growing season ("MAY-SEP" column), and the year ("ANNUAL" column). A valid daily mean is one for which 75% of the observations are available for each day, i.e., 18 hours. N is the number of days during each month, growing season, and year with at least 18 observations.
- (4) **Average Daily Mean.** The average of all valid daily means for the month, the growing season ("MAY-SEP" column), and the year ("ANNUAL" column). N is as in (3) above.
- (5) **Max Peak:Min Ratio.** The ratio of the Daily 1-Hr Maximum to the Daily 1-Hr Minimum. A ratio is computed only if a valid Daily Mean is computed and if the Daily 1-Hr Minimum is not equal to zero. N is the number of days with a valid Peak:Min ratio.
- (6) **Average Peak:Min Ratio.** The average of all Peak:Min ratios for the month, growing season, or year. N is as in (5) above.
- (7) **Max 9AM-4PM Average.** The maximum of all valid 9AM-4PM Averages computed for the month, growing season, or year. A valid 9AM-4PM Average is one which has 75% of the observations available during that time period (i.e., 6 hours. N is the number of days with valid averages.)

- (8) **Monthly 9AM-4PM Average.** The average of all valid 9AM-4PM Averages for the month, growing season, or year. N is as in (7) above.
- (9) **Max 7AM-7PM Average.** The maximum of all valid 7AM-7PM Averages computed for the month, growing season, or year. A valid 7AM-7PM Average is one which has 75% of the observations available during that time period, i.e., 9 hours. N is the number of days with valid averages.
- (10) **Monthly 7AM-7PM Average.** The average of all valid 7AM-7PM averages for the month, growing season, or year. N is as in (9) above.
- (11) **Monthly Mean.** The average of all 1-Hr ozone concentrations recorded during the month, growing season, or year. A mean is computed regardless of the number of hours with valid data. N is the number of hours with valid observations.
- (12) **SUM0 Exposure Index.** The monthly sum of all hourly ozone concentrations. Units are PPB-HR. The "MAY-SEP" column sums across the months of May through September to give the cumulative exposure for the growing season. The "ANNUAL" column sums across every month to give the cumulative exposure for the year. N is the number of hours with valid observations and is the same N as in (11) above.
- (13) **SUM60 Exposure Index.** The monthly sum of all hourly ozone concentrations equaling or exceeding 60 PPB. Units are PPB-HR. The "MAY-SEP" column sums across the months of May through September to give the cumulative exposure for the growing season. The "ANNUAL" column sums across every month to give the cumulative exposure for the year. N is the number of hours equaling or exceeding 60 PPB during the month, growing season, or year.
- (14) **SUM80 Exposure Index.** The monthly sum of all hourly ozone concentrations equaling or exceeding 80 PPB. Units are PPB-HR. The "MAY-SEP" column sums across the months of May through September to give the cumulative exposure for the growing season. The "ANNUAL" column sums across every month to give the cumulative exposure for the year. N is the number of hours equaling or exceeding 80 PPB during the month, growing season, or year.
- (15) **W126 Exposure Index.** The monthly sum of all hourly ozone concentrations where each concentration is weighted by a function that gives greater emphasis to the higher hourly concentrations while still including the lower ones. This weighting function provides a weighting value that is unique for each hourly ozone concentration. The weighting function, as described by Lefohn, Laurence, and Kohut¹ is:

$$w_i = \frac{1}{1 + 4403 \exp(-.126c_i)}$$

where

Weighting Function Used To Calculate W126 Exposure Index



w_i = weighting value for hourly concentration i ,
and
 c_i = hourly concentration i in PPB.

The graph of weighting value versus ozone concentration, in the figure to the left, illustrates the greater weights given to higher hourly ozone concentrations.

Each hour's weighting value is multiplied by its corresponding hourly concentration. This product is summed over all the valid hours in each month to calculate the monthly W126 exposure.

Thus, the monthly W126 exposure is:

$$W126 = \sum_{i=1}^n w_i c_i$$

where

- W126 = monthly W126 exposure index,
- w_i = weighting value for hourly concentration i ,
- c_i = hourly concentration i in PPB, and
- n = number of hours in the month with valid ozone concentrations.

The "MAY-SEP" column sums across the months of May through September to give the cumulative exposure for the growing season. The "ANNUAL" column sums across every month to give the cumulative exposure for the year. The exposure units are PPB-HR.

Because each hour contributes to this exposure index, N is the number of hours with valid observations and is the same N as in (11) and (12) above.

The U.S. Environmental Protection Agency usually considers air quality statistics, such as a mean, to be "valid" (i.e., representative of the parameter being estimated for the time interval in question) only if 75% or more of the total possible observations have been measured during that time interval. Therefore, one should exercise caution when comparing these statistics between months and sites, particularly those that are not averages (e.g., maxima and exposures) whenever the number of valid observations is less than 75% of the total possible.

References

1. Lefohn, A.S., J. A. Laurence, and R. J. Kohut. 1988. A Comparison of Indices That Describe the Relationship Between Exposure to Ozone and Reduction in the Yield of Agricultural Crops. *Atmospheric Environment* 22, 1229-1240.

4.2 AIR QUALITY GLOSSARY

Acid Deposition: Air pollution produced when acid chemicals are incorporated into rain, snow, fog, or mist.

Aerometric Information Retrieval System (AIRS): A computer-based database of U.S. air pollution information administered by the EPA Office of Air Quality Planning and Standards (U.S. Environmental Protection Agency).

AIRWeb: Air Resources Web, an air quality information retrieval system for U.S. parks and wildlife refuges developed by the Air Resources Division of the National Park Service and the Air Quality Branch of the Fish and Wildlife Service.

Air Pollutant: An unwanted chemical or other material found in the air.

Air Pollution: Degradation of air quality resulting from unwanted chemicals or other materials occurring in the air.

Air Quality: The properties and degree of purity of air to which people and natural and heritage resources are exposed (in the context of national parks).

Air Pollution Control Permitting Process: Process by which facilities are permitted to emit specified types and quantities of air pollutants.

Air Quality Related Values (AQRVs): Values including visibility, flora, fauna, cultural and historical resources, odor, soil, water, and virtually all resources that are dependent upon and affected by air quality. "These values include visibility and those scenic, cultural, biological, and recreation resources of an area that are affected by air quality." (43 Fed. Reg. 15016)

Ambient Air: Air that is accessible to the public.

Class I: Areas of the country set aside under the Clean Air Act to receive the most stringent degree of air quality protection.

Class II: Areas of the country protected under the Clean Air Act but identified for somewhat less stringent protection from air pollution damage than Class I, except in specified cases.

Clean Air Act: Originally passed in 1963, our current national air pollution control program is based on the 1970 version of the law. Substantial revisions were made by the 1990 Clean Air Act Amendments.

Continuous Sampling Device: An air analyzer that measures air quality components continuously.

Criteria: Information on health and/or environmental effects of pollution (in the context of criteria air pollutants).

Criteria Air Pollutant: A group of very common air pollutants regulated by EPA on the basis of criteria and for which a National Ambient Air Quality Standard is established (SO₂, NO₂, PM₁₀, Pb, CO, O₃).

Emissions: Release of pollutants into the air from a source.

Environmental Protection Agency (EPA): The federal agency responsible for regulating air quality.

Monitoring: Measurement of air pollution.

National Ambient Air Quality Standards (NAAQS): Permissible levels of criteria air pollutant established to protect public health and welfare.

Ozone (O₃): A criteria air pollutant that is a strong oxidizing agent, reactive with many other compounds and surfaces, and a health hazard in high concentrations. Ozone is formed by nitrogen oxides and organic compounds reacting in sunlight.

Source: Any place or object from which air pollutants are released. Sources that are fixed in space are stationary sources; sources that move are mobile sources.

Sulfur Dioxide (SO₂): A criteria air pollutant that is a gas produced by burning coal and some industrial processes.

* Recent updates to this glossary may be found on the NPSARD AIRWeb - <http://www.aqd.nps.gov/natnet/ard/glossary.htm>.

4.3 GLOSSARY OF AIR QUALITY UNITS

Units Conversion Table			
Parameter Type	Multiply	By	To Obtain
Pollutant	ppm	1000	ppb
	ppm	1960	$\mu\text{g}/\text{m}^3$ Ozone (at 25°C)
	ppm	2615	$\mu\text{g}/\text{m}^3$ Sulfur Dioxide (at 25°C)
	ppb	0.001	ppm
	ppb	1.960	$\mu\text{g}/\text{m}^3$ Ozone (at 25°C)
	ppb	2.615	$\mu\text{g}/\text{m}^3$ Sulfur Dioxide (at 25°C)
	$\mu\text{g}/\text{m}^3$ Ozone (25°C)	0.0005102	ppm
	$\mu\text{g}/\text{m}^3$ Ozone (25°C)	0.5102	ppb
	$\mu\text{g}/\text{m}^3$ Sulfur Dioxide (25°C)	0.0003824	ppm
	$\mu\text{g}/\text{m}^3$ Sulfur Dioxide (25°C)	0.3824	ppb
Wind Speed	m/s	2.05	mph
	mph	0.489	m/s
Solar Radiation	ly/min	697	w/m^2
	w/m^2	0.00143	ly/min
Precipitation	mm/hr	0.0394	in/hr
	in/hr	25.4	mm/hr
Temperature	$^{\circ}\text{C} + 17.78$	1.8	$^{\circ}\text{F}$
	$^{\circ}\text{F} - 32$	5/9	$^{\circ}\text{C}$
Where: ppm = parts per million ppb = parts per billion $\mu\text{g}/\text{m}^3$ = micrograms per cubic meter (at 25°C) m/s = meters per second mph = miles per hour ly/min = langley's per minute w/m^2 = watts per square meter mm/hr = millimeters per hour in/hr = inches per hour $^{\circ}\text{C}$ = degrees centigrade $^{\circ}\text{F}$ = degrees fahrenheit			