

**PETERSBURG NATIONAL BATTLEFIELD
GEOLOGIC RESOURCE MANAGEMENT ISSUES
SCOPING SUMMARY**

Trista L. Thornberry- Ehrlich
Colorado State University – Geologic Resource Evaluation
April 19, 2005



The entrance to the Union dug tunnel at Petersburg National Battlefield.
Photograph by Trista L. Thornberry- Ehrlich, 2005 (Colorado State University).

Table of Contents

Executive Summary	4
Introduction	5
Physiography	6
Geologic History of Central Virginia	7
Stratigraphy	10
Structure	12
Significant Geologic Resource Management Issues in Petersburg National Battlefield	13
Scoping Meeting Participants	21
References	22
Map of Petersburg National Battlefield.....	24

Executive Summary

A Geologic Resources Evaluation scoping meeting for Petersburg National Battlefield took place at Petersburg National Battlefield in Petersburg, VA on April 13, 2005. The scoping meeting participants identified the following list of geologic resource management issues. These topics are discussed in detail on pages 13- 20.

1. Shoreline changes at City Point and relative water level rises during storm surges. This is key to understanding erosion rates to protect the cultural resources.
2. Surface water quality and wetland management
3. Hydrogeologic characterization and groundwater quality to protect drinking supplies and understand how contaminants are moved through the system
4. Geologic hazards including radon and swelling soils with potential to influence new park land aquirements
5. Connections between geology and the Civil War history to appeal to visitors interested in a deeper connection to the landscape
6. Human impacts including gas pipelines, water lines, agricultural drainage ditches, and farm roads, nearby urban development
7. Mine- related features such as quarries, sand and gravel borrow pits
8. Sediment quality and sediment loading in park waterways with regards to sediment transport of contaminants and effects on riparian areas
9. Unique geologic features and Pre- Civil war historic artifacts
10. Future facilities development and characterization with regards to groundwater flow and other geologic parameters

Introduction

The National Park Service held a Geologic Resource Evaluation scoping meeting for Petersburg National Battlefield in Petersburg National Battlefield in Petersburg, Virginia on Wednesday, April 13, 2005. The purpose of the meeting was to discuss the status of geologic mapping in the park, the associated bibliography, and the geologic issues in the park. The products to be derived from the scoping meeting are: (1) Digitized geologic maps covering the park; (2) An updated and verified bibliography; (3) Scoping summary (this report); and (4) A Geologic Resource Evaluation Report which brings together all of these products.

Petersburg National Battlefield was designated a National Military Park on July 3, 1926 during Herbert Hoover's administration. It was later redesignated a National Battlefield on August 24, 1962 under John F. Kennedy. The park preserves part of the battlefield from the longest siege in American history. In 1864, Union forces dug in to cut-off Confederate supply lines to Richmond and Petersburg. Robert E. Lee's troops evacuated on April 2, 1865. The Civil War ended just days later. Petersburg National Battlefield covers 2,739 acres and is one of the larger military parks in the eastern United States. It protects a portion of the Harrison Creek watershed, not to mention the historical battle features including The Crater, earthworks, trenches, forts and, the Union dug tunnel.

Petersburg National Battlefield identified 8 quadrangles of interest, 9 with the possible addition of the DeWitt quadrangle. These quadrangles are all contained within the 30'x60' 1:100,000 Petersburg sheet. U.S. Geological Survey Bulletin 1567 covers portions of the Chester and all of the Hopewell quadrangles (1:24,000, 1987). The Virginia Department of Mineral Resources 1:500,000 scale 1993 state map has more detailed information relevant to Petersburg National Battlefield. Mapping in the area continues as part of the VDMR State Map project, though a definite time schedule has not been established.

Many other maps exist for the region that include coverage of the geology, oil and gas features, surficial geology, topography, groundwater features, land use, Landsat imagery, geochemical features, aeromagnetic- gravity, mineral and mineral potential, hazard features, stratigraphy, hydrogeology, structures, glacial features, karst features, etc. The maps are available from agencies such as the U.S. Geological Survey, the Virginia Division of Mineral Resources, the Geological Society of America, the American Geological Institute, the Maryland Geological Survey, the West Virginia Geological and Economic Survey, and the West Virginia Geological Survey. Additional mapping at a smaller scale within park boundaries will be more helpful for park resource management and interpretation.

Physiography

Petersburg National Battlefield Park lies within the transition between the Piedmont Plateau and Atlantic Coastal Plain physiographic provinces along the Fall Line. In the area of Petersburg, Virginia, the eastern United States is divided into 5 physiographic provinces with associated local subprovinces. These are, from east to west, the Atlantic Coastal Plain, the Piedmont Plateau, the Blue Ridge, the Valley and Ridge, and the Appalachian Plateaus provinces.

The eastward- sloping Piedmont Plateau is located between the Blue Ridge province along the eastern edge of the Appalachian Mountains and the Atlantic Coastal Plain province to the east. It formed through a combination of folds, faults, uplifts, and erosion. The resulting landscape of eastern gently rolling hills starting at 60 m (197 ft) in elevation becomes gradually steeper westward toward the western edge of the province and reaches 300 m (984 ft) above sea level. The Piedmont Plateau is composed of hard crystalline, igneous, and metamorphic rocks such as schists, phyllites, slates, gneisses, and gabbros. Soils in the Piedmont Plateau are highly weathered and generally well drained. The “Fall Line” or “Fall Zone” marks a transitional zone where the softer, less consolidated sedimentary rock of the Coastal Plain to the east intersects the harder, more resilient metamorphic rock to the west to form an area of ridges, waterfalls, and rapids.

The Atlantic Coastal Plain province is primarily flat terrain with elevations ranging from sea level to about 100 m (300 ft) in Maryland. Sediments eroding from the Appalachian Highland areas to the west formed the wedge- shaped sequence of soft sediments that were deposited intermittently during periods of higher sea level over the past 100 million years. These sediments are now more than 2,438 m (8,000 ft) thick at the Atlantic coast and are reworked by fluctuating sea levels and the continual erosive action of waves along the coastline. Large streams and rivers in the Coastal Plain province, including the James, York, Rappahannock, and Potomac, continue to transport sediment and to extend the coastal plain eastward. Beyond the province to the east is the submerged Continental Shelf province for another 121 km (75 miles).

The Blue Ridge contains the highest elevations in the Appalachian Mountain system. These are in Great Smoky Mountains National Park in North Carolina and Tennessee. Precambrian and Paleozoic igneous, sedimentary, and metamorphic rocks were uplifted during several orogenic events to form the core of the mountain range. Today this comprises the steep, rugged terrain now exposed after millions of years of erosion. Characteristics of the Blue Ridge province include steep terrain covered by thin, shallow soils, resulting in rapid runoff and low ground water recharge rates.

Geologic History of Central Virginia

Proterozoic Era – In the mid Proterozoic, during the Grenville orogeny, a supercontinent formed which included most of the continental crust in existence at that time. This included the crust of North America and Africa. The sedimentation, deformation, plutonism (the intrusion of igneous rocks), and volcanism associated with this event are manifested in the metamorphic gneisses in the core of the modern Blue Ridge Mountains (Harris et al., 1997). These rocks were deposited over a period of a 100 million years and are more than a billion years old, making them among the oldest rocks known from this region. They form a basement upon which all other rocks of the Appalachians were deposited (Southworth et al., 2001).

The late Proterozoic, roughly 600 million years ago, brought a tensional, rifting tectonic setting to the area. The supercontinent broke up and a sea basin formed that eventually became the Iapetus Ocean. In this tensional environment, flood basalts and other igneous rocks such as diabase and rhyolite added to the North American continent. These igneous rocks were intruded through cracks in the granitic gneisses of the Blue Ridge core and extruded onto the land surface during the break-up of the continental land mass (Southworth et al., 2001). Today these flood basalts comprise the Catoclin Greenstone. The Iapetus basin collected many of the sediments that would eventually form the Appalachian Mountains and Piedmont Plateau.

Early Paleozoic Era –

From Early Cambrian through Early Ordovician time orogenic activity along the eastern margin of the continent began again. The Taconic orogeny (~440- 420 Ma in the central Appalachians) was a volcanic arc – continent convergence. Oceanic crust, basin sediments, and the volcanic arc from the Iapetus basin were thrust onto the eastern edge of the North American continent. The Taconic orogeny involved the closing of the ocean, subduction of oceanic crust, the creation of volcanic arcs and the uplift of continental crust (Means, 1995). Initial metamorphism of the Catoclin Formation into metabasalts and metarhyolites, as well as the basin sediments (graywackes) into quartzites and phyllites occurred during this orogenic event.

In response to the overriding plate thrusting westward onto the continental margin of North America, the crust bowed downwards to the west creating a deep basin that filled with mud and sand eroded from the highlands to the east (Harris et al., 1997). This so-called Appalachian basin was centered on what is now West Virginia. These infilling sediments are today represented by the shale of the Ordovician (450 Ma) Martinsburg Formation (Southworth et al., 2001).

During the Late Ordovician, the oceanic sediments of the shrinking Iapetus Ocean were thrust westward onto other deepwater sediments of the western Piedmont. Sandstones, shales, siltstones, quartzites, and limestones were continuously deposited in the shallow marine to deltaic environment of the Appalachian basin. These rocks, now metamorphosed, currently underlie the Valley and Ridge province (Fisher, 1976).

This shallow marine to fluvial sedimentation continued for a period of about 200 My during the Ordovician, Silurian, Devonian, Mississippian, Pennsylvanian, and Permian Periods. This resulted in thick piles of sediments. The source of these sediments was from the highlands that were rising to the east during the Taconian orogeny (Ordovician), and the Acadian orogeny (Devonian).

The Acadian orogeny (~360 Ma) continued the mountain building of the Taconic orogeny as the African continent approached North America (Harris et al., 1997). Similar to the preceding Taconic orogeny, the Acadian event involved land mass collision, mountain building, and regional metamorphism (Means 1995). This event was focused further north than central Virginia.

Late Paleozoic Era – Following the Acadian orogenic event, the proto- Atlantic Iapetus Ocean was completely destroyed during the Late Paleozoic as the North American continent collided with the African continent. This formed the Appalachian mountain belt we see today and a supercontinent named Pangaea. This mountain building episode is called the Alleghanian orogeny (~325 – 265 Ma), the last major orogeny of the Appalachian evolution (Means, 1995). The deformation by folding and faulting produced the Sugarloaf Mountain anticlinorium and the Frederick Valley synclinorium in the western Piedmont, the Blue Ridge- South Mountain anticlinorium, and the numerous folds of the Valley and Ridge province (Southworth et al., 2001).

During this orogeny, rocks of the Great Valley, Blue Ridge, and Piedmont provinces were transported as a massive block (Blue Ridge – Piedmont thrust sheet) westward onto younger rocks of the Valley and Ridge along the North Mountain fault. The amount of compression was extreme. Estimates are of 20-50 percent shortening which translates into 125–350 km (75- 125 miles) of lateral translation (Harris et al., 1997).

Deformed rocks in the eastern Piedmont were also folded and faulted and existing thrust faults were reactivated as both strike slip and thrust faults during the Alleghanian orogenic events (Southworth et al., 2001). Paleoelevations of the Alleghanian Mountains are estimated at approximately 6,096 m (20,000 ft), analogous to the modern day Himalaya Range in Asia.

Mesozoic Era – Following the Alleghenian orogeny, during the late Triassic, a period of rifting began as the deformed rocks of the joined continents began to break apart from about 230- 200 Ma. The supercontinent Pangaea was segmented into roughly the continents that persist today. This episode of rifting or crustal fracturing initiated the formation of the current Atlantic Ocean and caused many block- fault basins to develop with accompanying volcanism (Harris et al., 1997; Southworth et al., 2001). These Mesozoic basins are scattered around the park area.

Thick deposits of unconsolidated gravel, sand, and silt were shed from the eroded mountains. These were deposited at the base of the mountains as alluvial fans and spread eastward to be part of the Atlantic Coastal Plain (Duffy and Whittecar 1991; Whittecar and Duffy, 2000; Southworth et al., 2001). The amount of material inferred from the now- exposed metamorphic rocks is immense. Many of the rocks exposed at the surface must have been at least 20 km (~10 miles) below the surface prior to regional uplift and erosion. The erosion continues today with the Potomac, Rappahannock, James, Rapidan, and Shenandoah Rivers, stripping the Coastal Plain sediments, lowering the mountains, and depositing alluvial terraces of the rivers, creating the present landscape.

Cenozoic Era – Since the breakup of Pangaea and the uplift of the Appalachian Mountains, the North American plate has continued to drift toward the west. The isostatic adjustments that uplifted the continent after the Alleghenian orogeny continued at a subdued rate throughout the Cenozoic Period (Harris et al., 1997). These adjustments may be responsible for occasional seismic events felt throughout the region.

Though glaciers from the Pleistocene Ice Ages never reached the central Virginia area (the southern terminus was in northeastern Pennsylvania), the intermittent colder climates of the ice ages played a role in the formation of the landscape at Petersburg National Battlefield. Freeze and thaw cycles at higher elevations led to increased erosion of large boulders and rocks by ice wedging. Sea level fluctuations during ice ages throughout the Pleistocene caused the baselevel of many of the area's rivers to change. During lowstands (sea level drops), the rivers would erode their channels exposing the deformed bedrock of the Piedmont Plateau. During oceanic highstands, the river basins flooded and deposition resulted in deposits of beach sediments around and beneath the park.

Stratigraphy

Petersburg National Battlefield sits on the eastern edge of the Fall Line, the boundary between the Atlantic Coastal Plain and Piedmont Plateau physiographic provinces. The term Fall Line, is a bit of a misnomer since the boundary is actually a zone of fluctuating width along the eastern edge of the Piedmont. The Fall Zone is the term used herein to describe the transition from unconsolidated Coastal Plain deposits to hard, metamorphosed rocks of the Piedmont. Piedmont rocks are present at great depths below the thick, wedge-shaped deposits. In the Petersburg area, the Fall Zone is where rivers such as the James that rise and fall with tides hit zones of rapids and waterfalls.

Bedrock exposures within the park are rare. The oldest rocks in the Petersburg National Battlefield are exposed along the deepest eroded reaches of the area's rivers. The ~300 Ma Petersburg Granite is probably the oldest unit within the park area (Clark, meeting communication, 2005). This rock is comprised of foliated to non-foliated, granite to granodiorite containing lenses of biotite gneiss and amphibolite (Berquist, 2003). To the west of the park and beneath the coastal plain sediments are granitic gneisses and diorite gneiss as well as several intrusive rocks including granite, quartz monzonite, and granodiorite. These gneissic rocks are Late Proterozoic to Early Paleozoic in age (Mixon et al., 2000). Intermixed with these intrusive rocks and becoming more predominant further south are the metamorphic rocks of the eastern Piedmont including biotite gneiss, mica schist, metabasalts, and phyllonite (VDMR, 1993)

Several Mesozoic age extension basins including the Richmond Mesozoic Basin exist around the battlefield area to the west. These basins formed during crustal extension accompanying the formation of the Atlantic Ocean. These basins were typically bounded by linear normal faults. The basins flooded and filled with sediments and were occasionally intruded by diabase dikes. The Richmond Mesozoic Basin contains dinosaur footprints in sediments more than 762 m (2,500 ft) thick on the western side. These sediments include sands, gravels, silts, and muds eroded from the Appalachian highlands.

Surrounding the battlefield units of the park are younger deposits of Atlantic Coastal Plain rocks. The Lower Cretaceous Potomac Formation is exposed by erosion along the Appomattox River. This formation is comprised of feldspathic quartz sand and sandstone, silty channel-bar deposits, and lignitic sandy silt and clay layers. Atop the Potomac Formation is the upper Paleocene Aquia Formation of glauconitic sands, silts, clays, and containing some scant fossil layers. Nanjemoy Formation deposits from the lower Eocene overlie the Aquia Formation. These are glauconitic sands, clays, silts, and mixed layers. This unit is as much as 64 m (210 ft) thick. Miocene age sands and gravels as much as 10 m (33

ft) thick overlie the Aquia and Nanjemoy Formations in the park area (Dischinger, 1987; Mixon et al., 2000).

The upper Miocene Eastover Formation is exposed by erosion along local rivers. This formation is comprised of fine quartzose sand layers interbedded with clayey silts. Union forces dug tunnels through this unit to breach Confederate lines and to plant explosives beneath a fort near Elliot's Salient. Younger deposits include the Yorktown Formation and Pliocene sands and gravels. The Yorktown Formation is a maximum of 25 m (82 ft) thick and contains quartz and feldspar sands mixed with lesser clays and silts. Beach sediments within these two units attest to glacially induced marine highstands where the shoreline was near present day Richmond and Petersburg 4 Ma (Berquist, meeting communication, 2005). These beach sediments contain heavy minerals, including titanium, sparking mining interest to the southeast in Dinwiddie County.

The upper Pliocene Bacons Castle Formation includes gravelly sand and sandy-silty- clayey upper layers. These are often found in high- level terrace areas. The more recent deposits at the park include various Quaternary age units. The fine to coarse sand, gravel, silt and clay of the Windsor Formation is of lower Pleistocene to upper Pliocene age. The gravelly Charles City Formation, the sands and silts of the Chuckatuck Formation as well as the Shirley Formation of coarse sands, gravels, pebbles and occasional boulders are of the middle Pleistocene (Mixon et al., 2000). The Shirley Formation deposit has basal units of sand and gravel, grading to muds in the upper beds. This 200 Ka unit is also from an interglacial highstand (Berquist, meeting communication, 2005). The upper Pleistocene Tabb Formation contains sands, gravels, silts and clays and has three members: the Poquoson, the Lynnhaven, and the Sedgefield Members (Mixon et al., 2000).

The youngest deposits at Petersburg National Battlefield include thick alluvium deposits of sand, gravel, silt and clays as much as 15 m (49 ft) thick, marsh and swamp deposits along larger rivers, shelly sands, and artificial fill from construction of roads, dams, bridges, landfills, and highways.

Structure

The Richmond Mesozoic Basin dominates the structural features present west of Petersburg National Battlefield. This feature is bounded by high angle normal faults. To the west, across a broad zone of sheared mylonite and phyllonite are Middle Proterozoic metamorphic gneisses of the Goochland Terrane. To the east of the basin are intrusive rocks, including granites, quartz monzonite, and granodiorite (VDMR, 1993). Another shear zone penetrates the eastern Piedmont metamorphic rocks south of the battlefield. This zone contains highly deformed mylonites and phyllonites.

Faults within the park area have had a pronounced influence on landform development. Most of these are not well exposed. Several large, parallel, low- to high- angle thrust faults dipping to the southeast are present far to the west trending northeast- southwest. These thrust faults moved large masses of Piedmont rocks and associated plutons atop younger rocks to the west. Many of these older faults were brittlely reactivated during crustal extension accompanying the opening of the Atlantic Ocean. Most of these faults have not been mapped due to lack of exposure, but most rock quarries in the area have one or more large scale faults (Berquist, meeting communication, 2005).

Closer to the park are several smaller scale faults surrounding the Mesozoic basin and further east into the Coastal Plain sediments. High- angle reverse faults are present along the eastern side of the Piedmont. Faulting within the Coastal Plain, though relatively small in scale is significant because it reveals very recent tectonic activity in the area. A north- trending zone of reverse faults was mapped by Dischinger (1987) as the Dutch Gap fault zone. This zone extends at least 13 km from the northwestern corner of the battlefield northward across the James River. In measurements of offset on the contact between the Cretaceous Potomac Formation and the Paleocene Aquia Formation, up to 20 m (* ft) of displacement is observed.

Highpoints near Bailey Creek and the southward reach of the James River from Richmond to Drewrys Bluff indicate the possible presence of other faults, parallel to the Dutch Gap fault zone. An increase in sediment thickness north of the James River may indicate the presence of a fault downdropped block to the north that would be perpendicular to the Dutch Gap system (Dischinger, 1987).

Significant Geologic Resource Management Issues at Petersburg National Battlefield

i. Shoreline changes and water level rise

The water level of the James River, in the vicinity of the City Point unit in Hopewell, VA, is largely controlled by tidal fluctuations coming up the Chesapeake Bay. In addition to regular tidal fluctuations, storm surges can dramatically increase the relative water level to flooding conditions.

One of the major goals of the park is to present the historical context of the area; this includes preserving and restoring any old buildings and the landscape around them. Maintaining this battle landscape often means resisting natural geologic changes, which presents several management challenges. Geologic slope processes such as landsliding, slumping, chemical weathering, and slope creep are constantly changing the landscape at the park.

The river and other local streams are changing position constantly as part of natural meandering river flow. These shoreline changes threaten existing park facilities and the historical context of the landscape. Large blocks of rock are sloughing off the bluffs at the City Point unit where the James and Appomattox Rivers converge. Storm events including hurricanes and tropical storms, such as Hurricane Isabel in 2003 and Tropical Storm Ivan in 2004, increase this effect with rapidly rising, erosive waves. Atop the bluffs at City Point are a 1763 historic home (Eppes house), Grant's cabin, and several archaeological sites.

Research and monitoring questions and suggestions include:

- Identify areas prone to slope failure during intense storm events.
- Investigate remedial techniques to reduce sloughing at City Point.
- Monitor erosion rates and shoreline changes along the James River and compare to previous conditions using aerial photographs where available.
- What are the effects of increased erosion on aquatic ecosystems at the park?
- Is runoff in the park increasing due to surrounding development? If so, are there any remedial efforts the park can undertake to reduce this impact?
- Should the park preserve (recreate) historic landscapes at the expense of natural processes?
- Should the park target certain areas for restoration and leave others to natural processes?
- Develop an interpretive program detailing the balance between cultural context and natural processes at Petersburg National Battlefield.

2. Surface water quality and wetland management

Many streams cross the landscape within and surrounding Petersburg National Battlefield. These include Harrison's Creek, Rohoic Creek, Poor Creek, Hatcher Run (and accompanying 4- 5 acre pond), Lieutenant Run, Brickhouse Run, Cattail Run, Arthur Swamp, Second Swamp, Picture Branch, Rocky Branch, Wallace Creek, Gravelly Run, Reedy Branch, and Bailey Creek, in addition to the large rivers, the James and the Appomattox. Because of this, the quality of the surface water at the park is very important to park management and by extension, the surrounding communities. Flooding and channel erosion are naturally occurring along most of the streams and rivers within the park. Poor Creek is cutting deep channels near The Crater, causing some slumping along the banks. Interesting views of sediment layers are also exposed through this process and may yield cultural and geological interpretation opportunities.

Several riparian wetlands exist within the park and are threatened by increased flow and floods, as well as by sediment loading. These have been inventoried by the National Wetlands Inventory program. Though small in scale, wetlands are typically considered indicators of overall ecosystem health and should be researched and monitored periodically. Surface water quality at the park is threatened by ground compaction due to increased use as well as increases in impervious surfaces such as parking lots and roadways. These features increase the amounts of seasonal runoff as sheet flow.

Alterations to park vegetation along the steep, exposed slopes lead to changes in the hydrologic regime at the park. For example, clearing of trees and their stabilizing roots for historical restoration, can lead to increased erosion, contribute to slumps and landslides, and thereby increase sediment load in nearby streams.

Research and monitoring questions and suggestions include:

- Monitor water and soil quality in wetlands to establish as basis for comparison of future conditions.
- Use aerial photographs to study changes to wetland distribution through time.
- Accurately map park use areas and target streams and areas of erosion for remediation.
- Determine any hotspots for water contamination. Remediate and monitor results.
- Research planting new vegetation along vulnerable reaches of park streams to prevent excess erosion and sediment loading.

3. Hydrogeologic characterization and groundwater quality

The resource management needs to understand how water is moving through the hydrogeologic system into, under, and from the park. This is critical to understanding the impacts of human induced contaminants on the ecosystem. The interaction between groundwater flow and the overall water quality should be quantitatively determined at the park. Little data regarding the nature of the hydrogeologic system at the park exists. Park resource personnel also need to understand how the water table might change over time to manage water resources. Municipal water and a well at Poplar Grove are the primary drinking water supply sources for the park. There are several wells throughout the park that could be used for monitoring of ground water quality. It would be useful to perform tracer studies in these wells to see how quickly and in what direction water is moving through the system.

Research and monitoring questions and suggestions include:

- Inventory groundwater levels at all park units.
- Test for and monitor phosphate and volatile hydrocarbon levels in the groundwater at the park, focusing on areas near facilities.
- Inventory and map any existing springs in the park, especially with regards to their potential historical importance.
- Test water quality at any existing springs in the park.
- Create hydrogeologic models for the park to better manage the groundwater resource and predict the system's response to contamination.

4 Geologic hazards

Radon is a naturally occurring radioactive gas, derived from radioactive decay of radium that is a product of uranium radioactive decay. Both radium and uranium are present in at least trace amounts in almost any kind of rock or soil. Particular rock types are more prone to high uranium concentrations including granites, shales, and phosphates. Radon is tasteless, odorless, and colorless. According to the Environmental Protection Agency (EPA) there is no safe radon level. Any exposure to the gas has serious associated health risks.

In wells tested near the park boundaries to the southwest, radon was detected. Dinwiddie County has high radon concentrations in the soils. This issue must be addressed when potential new park lands are assessed.

Swelling soils occur in the Chesterfield County area, part of which is in the western quadrangles of interest for Petersburg National Battlefield. Swelling soils (clays) expand when water saturated and shrink when dry. In extreme cases, these soils can undermine roads, buildings, and other structures. In less extreme ways, they can affect trails and other visitor use facilities.

Research and monitoring questions and suggestions include:

- Should radon be tested at wells in all park units?
- Use soil surveys and geologic maps to determine areas especially prone to radon problems.
- Map locations of high swelling clay concentrations to avoid for future development.

5. Connections between geology and the Civil War history

In many Civil War battles, it was the advantage that familiarity with terrain, preparation to utilize of the natural features of the area, and the manipulation of the focal points, the gaps, ravines, cuts, hills, and ridges gave to one side or another that was to decide the outcome. The rolling hills and gentle landscape and topography at Petersburg National Battlefield are defined by the geology and hydrology. This setting dictated the placement of the towns, strategy and encampment of troops, escape routes, river crossings, railroads, and the development of outlying areas including City Point, Taylor farm, various forts within the siege area, and the earthworks themselves. Union forces dug tunnels (the location of only one of which is known) through the substrate, often in the unconsolidated Eastover Formation. The most famous tunnel resulted in The Crater amidst the Confederate line near Elliot's Salient. In addition to influencing battles, the landscape and topography also affected how troops and supplies were transported during the Civil War.

Runoff erodes sediments from any open areas and carries them down streams and gullies. Erosion naturally diminishes higher areas such as ridges and hills, foundations, earthworks, degrades bridge foundations, erodes streams back into restoration areas, and fills in the lower areas such as trenches, railroad cuts and stream ravines distorting the historical context of the landscape. Earthworks at the park are small. These 140- year- old features are eroding away.

Interpreters make the landscape come alive for visitors and give the scenery a deeper meaning. Because geology forms the basis of the entire ecosystem, it is directly responsible for the history at Petersburg National Battlefield. Thus, geologic features and processes should be emphasized to improve the visitor's experience. The website for the park needs to be updated for geologic content and connections with other scientific and cultural disciplines.

Research and monitoring questions and suggestions include:

- Create interpretive programs concerning geologic features and processes and their effects on the battleground history of the park.
- Encourage the interaction between geologists and the interpretive staff to come up with a list of features and programs to execute.

- Create a general interest map with simple explanatory text on geologic battle and settlement influences for visitors to the park.
- Update the park website relating geology with other resources.
- Perform surveys to discover reported Union tunnels in other areas of the park.

6. Human impacts

The area surrounding Petersburg National Battlefield is becoming increasingly populated. As development continues, conservation of any existing forest-meadow community types becomes a critical concern. Understanding the geology beneath the biotic communities becomes vital to their management. Park management of the landscape for historic preservation purposes compliments the preservation of these forests.

Humans began settling the Petersburg area in the late 1600's to early 1700's, stemming from the earliest settlements along the James River at Jamestown in 1607. Their farming and homestead activities created an unnatural landscape that persists today at Petersburg National Battlefield. Minor irrigation features, removal of soil and rocks, grazed pastures, and homestead foundations dot the landscape.

Human impacts continue today as water lines, gas lines, power lines, roads, and buildings increase on the landscape. The gas and water lines running under the park are buried to a depth of only 1.5 m (5 ft). The gas line feeds the entire Tidewater area of Virginia. Additionally, trails, visitor use areas, imported (invasive species), acid rain, and air and water pollution take their toll on the landscape. Resource management of these impacts is an ongoing process.

Research and monitoring questions and suggestions include:

- Perform acid rain measurements and correlate with the underlying bedrock to determine if any buffering effects occur. Relate this information to the water quality for the park.
- Should the unnatural landscapes created by early settlers be remediated?
- Are soils becoming more acidic due to acid rain?
- Monitor chemical alterations in bedrock.
- Cooperate with local developers to minimize impact near park areas.
- Consult conservation groups regarding cooperative efforts to increase the areas of relevant parklands and protect more of the region around the battlefields from development.
- Promote environmentally sound methods of developing land parcels including partial clearing of trees and proper construction of stable slopes.

7. Mine- related features

Small- scale abandoned sand and gravel pits dot the landscape within the units at Petersburg National Battlefield. The park filled many of these features. New lands being considered for park acquisition are near a sand quarry and a Petersburg Granite quarry. Petersburg Granite has long been quarried as a building material and many of the streets and buildings in surrounding towns were built from it. The state of Virginia has an inventory of mine features such as small open pit mines. Mines are constantly being identified and scanned into their database.

These features pose several concerns for resource management. Visitor safety is a constant concern wherever open pits occur. Most mine features, accessible to the public have been blocked or filled.

Research and monitoring questions and suggestions include:

- Comprehensively inventory mine- related features at Petersburg National Battlefield, consulting aerial photographs and historic records if necessary.
- Consult the Virginia Department of Mineral Resources database for abandoned mine information.
- Perform a stability survey, focusing on any open gravel and sand pits.

8. Sediment quality and sediment loading

Erosion of the landscape at Petersburg leads to increases in sediment being carried by the area's rivers. Sediment loads and distribution affect aquatic and riparian ecosystems, and sediment loading can result in changes to channel morphology and increase overbank flooding frequency.

Fine- grained sediments are vital in the overall fluvial transport of contaminants in a water system. Channel storage of fine sediment (and the contaminants contained therein) follows a seasonal cycle. This cycle is subject to not only human disturbance, but hydrologic variability with increased availability during the high stands of spring and decreased availability during the low stands of autumn (Miller et al., 1984). Nearby chemical plants and paper mills are introducing contaminants into the James and Appomattox Rivers. Dredging proposals for tour boats on the Appomattox River were considered too risky because of the contaminants they might disturb in the river bottom sediments (Blumenshine and Shockley, meeting communication, 2005).

Fine- grained sediments do not travel downstream in a single pulse, but are often resuspended bottom material (Miller et al., 1984). This intermittent transport of contaminants and fine- grained sediment increases the affected area. This could

pose a threat to the park if a focused, severe contamination were to occur within a watershed such as Harrison Creek or the Appomattox River..

Research and monitoring questions and suggestions include:

- Monitor fine- sediment load in Harrison Creek. Appomattox River?
- Cooperate with local agencies to determine amounts, types, and durations of exposure of the watershed to contaminants.
- Research stream sediment loads and qualities to determine their effects on aquatic and riparian biota.
- Is sediment loading in the park streams following a seasonal pattern?

9. Unique geologic features

The 300 Ma Petersburg Granite is exposed in the deepest eroded areas around Petersburg National Battlefield. Within the granite are pegmatite veins (exceptionally coarse- grained igneous rock) containing crystals of striped beryl, zircons, and pyrite. The type section for this rock unit is in the area. For geologists, a type section is the locality where the definition and description of the unit are established. This has important implications for future geologic research. In addition, large boulders, some the size of cars, are located in fields at Petersburg and along Highway 85. These unique features may represent glacial erratics, or pieces of rock that floated away from their source on rafts of ice that are dropped when the ice melts, often far from their origin. During interglacial periods, sea level rises as more water is freed from glacial ice. In eastern Virginia, beach deposits far west of the current coastline are areas of heavy metal concentrations. These concentrations have sparked interest in titanium mining in Dinwiddie County.

Research and monitoring questions and suggestions include:

- Are the large boulders found on the landscape at Petersburg glacial erratics?
- Where were these boulders derived from?
- Educate the public about the presence of a type section in the park area.
- Develop an interpretive program regarding the mineral resources in the area including the Petersburg Granite, crystals within the pegmatites, and titanium and other heavy minerals found in nearby beach deposits.

10. Future facilities

Many geologic factors must be taken into account during decision making regarding siting future facilities. Lands in Dinwiddie County are being considered for acquirement by the park. The presence of radon could affect the development of buildings and wells. Similarly, high concentrations of shrink and swell clays could undermine any structure. Knowing the locations of springs and

understanding the hydrogeologic system helps avoid problems with groundwater flow. Waste facilities should be sited taking hydrogeology into account as well as an understanding of the substrate.

Research and monitoring questions and suggestions include:

- Consult local geologic experts when planning future facilities at the park.
- Locate areas of high swelling clay concentrations, springs, groundwater flow, high slope, unstable substrate, etc. to avoid for future facilities.

Scoping Meeting Participants

NAME	AFFILIATION	PHONE	EMAIL
Kristen Allen	NPS- RICH	804- 795- 5019	kristen_allen@nps.gov
Rick Berquist	Virginia Division of Mineral Resources	757- 221- 2448	rick.berquist@dmme.virginia.gov AND crberq@wm.edu
Tim Blumenshine	NPS- PETE	804- 732- 0170, ext. 303	tim_blumenschine@nps.gov
Mark Carter	AASG- Virginia Division of Mineral Resources	434- 951- 6357	mark.carter@dmme.virginia.gov
Jim Comiskey	NPS, MIDN	540- 654- 5328	jim_comiskey@nps.gov
Tim Connors	NPS, Geologic Resources Division	303- 969- 2093	tim_connors@nps.gov
Nate Irwin	NPS- PETE	804- 862- 3019, ext. 301	nathaniel_irwin@nps.gov
Gregg Kneipp	NPS- FRSP	540- 654- 5331	gregg_kneipp@nps.gov
Stephanie O'Meara	NPS, Natural Resources Information Division	970- 225- 3584	Stephanie_O'Meara@partner.nps.gov
Dave Shockley	NPS- PETE	804- 732- 0171, ext. 305	dave_shockley@nps.gov
Trista Thornberry-Ehrlich	Colorado State University	757- 222- 7639	tthorn@cnr.colostate.edu

References

- Berquist, C.R., Jr., 2003, Digital Representation of the 1993 Geologic Map of Virginia – Expanded Explanation. Commonwealth of Virginia Department of Mines, Minerals, and Energy, Division of Mineral Resources, Publication 174, 89 p.
- Commonwealth of Virginia Department of Mines, Minerals, and Energy, Division of Mineral Resources (VDMR), 1993, Geologic Map of Virginia, 1:500,000 scale.
- Condie, K.C., Sloan, R.E., 1998, Origin and Evolution of the Earth, Principles of Historical Geology. Prentice- Hall, Inc., 498 p.
- Dischinger, J.B., 1987, Late Mesozoic and Cenozoic stratigraphic and structural framework near Hopewell, Virginia. U.S. Geological Survey, Bulletin 1567, 1:24,000 scale.
- Duffy, D.F., Whittecar, G.R., 1991, Geomorphic development of segmented alluvial fans in the Shenandoah Valley, Stuarts Draft, Virginia. Abstracts with Programs - Geological Society of America, vol.23, no.1, p.24.
- Fisher, G.W., 1976, The geologic evolution of the northeastern Piedmont of the Appalachians. Abstracts with Programs - Geological Society of America, vol.8, no.2, p.172- 173.
- Harris, A.G., Tuttle, E., Tuttle, S.D., 1997, Geology of National Parks. Kendall/Hunt Publishing Company, 759 p.
- Means, J., 1995, Maryland's Catoctin Mountain parks; an interpretive guide to Catoctin Mountain Park and Cunningham Falls State Park. McDonald & Woodward Publishing Company : Blacksburg, VA, United States, 168 p.
- Miller, A.J.; Smith, J.A.; Shoemaker, L.L., 1984, Channel storage of fine- grained sediment in the Monocacy River basin., Eos, Transactions, American Geophysical Union, Vol. 65, Issue 45, pp. 888
- Mixon, R.B., Pavlides, L., Powars, D.S., Froelich, A.J., Weems, R.E., Schindler, J.S., Newell, W.L., Edwards, L.E., Ward, L.W., 2000, Geologic Map of the Fredericksburg 30'x60' Quadrangle, Virginia and Maryland. U.S. Geological Survey, Geologic Investigations Series Map I- 2607, 1:100,000 scale.

- Nickelsen, R.P., 1956, Geology of the Blue Ridge near Harpers Ferry, West Virginia. Geological Society of America Bulletin, vol.67, no.3, p.239- 269.
- Onasch, C.M., 1986, Structural and metamorphic evolution of a portion of the Blue Ridge in Maryland. Southeastern Geology, Vol. 26, Issue 4, pp. 229- 238.
- Southworth, S., Brezinski, D.K., Orndorff, R.C., Chirico, P.G., Lagueux, K.M, 2001, Geology of the Chesapeake and Ohio Canal National Historical Park and Potomac River Corridor, District of Columbia, Maryland, West Virginia, and Virginia; A, Geologic map and GIS files (disc 1); B, Geologic report and figures (disc 2). U. S. Geological Survey, Open- File Report: OF 01- 0188.
- Thomas, W.A., Chowns, T.M., Daniels, D.L., Neatherly, T,L., Glover, L., Gleason, R.J., 1989, The subsurface Appalachians beneath the Atlantic and Gulf coastal plains. The Geology of North America, vol. F- 2, Geological Society of America.
- Whittecar, G. R., Duffy, D.F., 2000, Geomorphology and stratigraphy of late Cenozoic alluvial fans, Augusta County, Virginia, U.S.A. In: Regolith in the Central and Southern Appalachians, Clark, G.M., Mills, H.H., Kite, J.S., eds., Southeastern Geology, vol.39, no.3- 4, p.259- 279.

Map of Petersburg National Battlefield

(Eastern portion)



